

# How brains of doers differ from those of procrastinators

August 22 2018

---



Two areas of the brain apparently determine if an individual tends to put tasks off or to tackle them straightaway. Credit: RUB, Marquard

Researchers at Ruhr-Universität Bochum have analysed why certain people tend to put tasks off rather than tackling them directly. Using

magnetic resonance imaging (MRI), they identified two brain areas whose volume and functional connectivity are linked to an individual's ability to control their actions. The research team headed by Caroline Schlüter, Dr. Marlies Pinnow, Professor Onur Güntürkün, and Dr. Erhan Genç from the Department of Biopsychology published the results in the journal *Psychological Science* on 17 August 2018.

## Two areas of the brain linked to action control

The biopsychologists examined 264 women and men in an MRI scanner. They assessed the volume of individual brain regions and the [functional connectivity](#) between them. In addition, all participants completed a survey measuring their own ability to execute action control.

Individuals with poor action control had a larger [amygdala](#). Moreover, the functional connection between the amygdala and the so-called dorsal anterior cingulate cortex (dorsal ACC) was less pronounced. "These two areas of the brain had already been linked with action control in former studies," says Erhan Genç.

## Assessing and selecting actions

The primary function of the amygdala is to assess different situations with regard to their respective outcomes and to warn us about potential [negative consequences](#) of particular actions. The dorsal ACC uses these information in order to select actions that are to be put into practice. Moreover, by suppressing competing actions and emotions, it ensures that the selected action can be successfully completed.

If the interplay between amygdala and dorsal ACC is impaired, action control can no longer be successfully executed, according to the theory put forward by the researchers. "Individuals with a higher amygdala

volume may be more anxious about the negative consequences of an action – they tend to hesitate and put off things," speculates Erhan Genç. "Due to a low functional connection between amygdala and dorsal ACC, this effect may be augmented, as interfering negative emotions and alternative actions might not be sufficiently regulated."

## Learnable or not?

Future studies will have to show if the degree of action control can be modified through specific training or brain stimulation. "Even though the differences regarding our ability to control our actions affect our private and professional success as well as our mental and physical health to a considerable degree, their neural foundations haven't as yet been sufficiently studied," says Caroline Schlüter, who addresses this issue in her Ph.D. thesis.

**More information:** Caroline Schlüter et al. The Structural and Functional Signature of Action Control, *Psychological Science* (2018). [DOI: 10.1177/0956797618779380](https://doi.org/10.1177/0956797618779380)

Provided by Ruhr-Universität-Bochum

Citation: How brains of doers differ from those of procrastinators (2018, August 22) retrieved 24 April 2024 from <https://medicalxpress.com/news/2018-08-brains-doers-differ-procrastinators.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--