

## **Delay eating breakfast and eat dinner early if you want to lose body fat – new study**

August 30 2018, by Jonathan Johnston And Rona Antoni

---



Credit: Nork Photography from Pexels

Time-restricted eating (also called time-restricted feeding) is a new

dietary concept that involves reducing the time between the first and last calorie consumed each day. There is strong evidence to support the health benefits of time-restricted eating (TRE) in animals, and recent small studies by our research group and others suggest possible benefits for humans, too.

A lot of [recent research in animals and humans](#) has shown clear links between our circadian (daily) rhythms, metabolism and nutrition. Circadian clocks exist throughout our bodies, including in tissues that have a big influence on our metabolic health.

Broadly speaking, these clocks enable our bodies to efficiently deal with food intake when we are awake and release energy from stores when we are asleep. A consequence of this is that we process food consumed during the day more efficiently than an identical meal consumed during the evening or at night.

There is, indeed, increasing evidence supporting the idea that our [metabolic health](#) is not just regulated by what we eat, but also when we eat. This interaction between circadian rhythms and nutrition is now often referred to as "[chrononutrition](#)".

One of the ways that chrononutrition is being applied to dietary health is with TRE. [Animal studies](#) in species as diverse as mice and fruit flies have demonstrated that TRE can reduce obesity caused by a high-fat diet and reduce the risk of heart and metabolic disease. Metabolic and nutritional studies in animals don't always apply to humans, however, so it is important to test TRE in humans before health claims can be made.

## **TRE in humans**

An early indication of the effects of TRE in humans is available in the form of Ramadan studies. Islamic religious fasting restricts eating and

drinking to nighttime, between sunrise and sunset. Recent [meta-analyses of these observational Ramadan studies](#) show improvements in some risk markers for diabetes and heart disease.

These studies are difficult to interpret, however, because Ramadan involves a change from daytime to nighttime eating, as well as changes in meal duration. Changes also tend to be short-lived and limited to the one-month fasting period.

In human intervention studies, TRE has been achieved in several ways, including [avoiding eating at night](#), restricting intake to a fixed [four](#), [six](#) or [eight](#) hour eating window, or by symmetrically reducing the eating window by three hours, relative to normal eating patterns.



Credit: Nork Photography from Pexels

The latter is an approach we used in a recent pilot study, published in [the Journal of Nutritional Science](#). We split participants into two groups. One group – the TRE group – delayed their breakfast by 90 minutes and had their dinner 90 minutes earlier. The other group (the control group) ate meals as they would normally.

Participants had to provide blood samples and complete diet diaries before and during the ten-week intervention and complete a feedback questionnaire immediately after the study.

We found statistically significant reductions in calorie intake and body fat in the TRE group, compared with controls. Our data also suggested that TRE might benefit blood markers of metabolic disease, including fasting blood sugar (glucose) concentration. Questionnaire data indicated that some, but not all, TRE participants felt the eating pattern was one they could continue with.

It should be noted that pilot studies are designed to generate preliminary data and help with the design of future large-scale research. The number of participants in pilot studies is small, so no definitive conclusions can be drawn from the results.

## **Early days**

In general, TRE studies have produced mixed results, but this should not be surprising, given the diversity of protocols. Various forms of TRE have been shown to either improve or have little effect on risk factors

for heart disease and diabetes. Some studies have shown improvements in body composition, such as reduced body fat, in both overweight and [obese](#) people, and also in [men who regularly exercise](#), although this is not found across all studies.

One common finding is that people tend to reduce [food intake](#) during TRE, even when asked not to. This is itself a finding that could have major health benefits. But even where energy restriction is absent or small, evidence suggests that [TRE can still influence metabolism](#), meaning that the benefits of TRE may go beyond simply regulating energy intake.

Important issues that still need to be investigated include whether some feeding-fasting times work better than others, how TRE might benefit health even when [calorie intake](#) is the same, and whether TRE benefits patients with [metabolic disease](#), such as type 2 diabetes.

In this fast-moving field, we and others are now looking to expand upon the initial results described here. Even when living in our modern 24-hour society, there may be important [health benefits](#) from eating in a pattern that fits with our internal [circadian rhythms](#).

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Delay eating breakfast and eat dinner early if you want to lose body fat – new study (2018, August 30) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-08-breakfast-dinner-early-body-fat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.