

Breast cancer surgery in frail elderly women linked to poor results

August 29 2018



Credit: CC0 Public Domain

An analysis of more than a decade of U.S. nursing home data has shown that breast cancer surgery is associated with high rates of mortality and hospital readmission, along with loss of functional independence, for

frail nursing home residents.

In a study appearing Aug. 29, 2018, in *JAMA Surgery*, UCSF researchers found that 58 percent of women who resided in a nursing home for more than 90 days before [breast cancer surgery](#) experienced significant functional decline one year after surgery. The study found that women with functional impairment in their daily activities prior to treatment had the highest rates of one-year mortality and functional decline. Patients with prior cognitive impairment also had higher rates of functional decline after one year.

"Surgery often cures the cancer, but can have a negative impact on elderly patients' everyday activities and worsen their quality of life," said lead author Victoria Tang, MD, MAS, assistant professor of geriatrics and of hospital medicine at UCSF and the affiliated San Francisco VA Health Care System. "This study shows that for frail, elderly patients, breast cancer care should be individualized and goal-oriented, with the option of only providing hormonal therapy or symptom management, instead of surgery."

Breast cancer surgery is the most common cancer operation performed in nursing home residents, constituting 61 percent of procedures. More than half of female nursing home residents are identified with suspected breast cancer through screening or physical exam, and about two-thirds of those are referred for diagnosis or treatment.

In the *JAMA Surgery* study, Tang and her colleagues used 2003-2013 claims from all U.S. Medicare nursing homes to review data for 5,969 women ages 67 and older who had lived in a nursing home for at least 90 days and who underwent inpatient breast cancer surgery. In this group (83 percent white, 57 percent cognitively impaired), 61 percent (3,661) of the patients received the most invasive treatment, known as axillary lymph node dissection with lumpectomy or mastectomy (ALND).

Another 28 percent (1,642) received a mastectomy, and 11 percent (666) underwent the least invasive lumpectomy. Researchers examined 30-day and one-year mortality, hospital readmission rates, and functional status in activities of daily living, such as eating, dressing and using the bathroom.

The researchers noted that the rates were high for 30-day readmission (16 percent overall, 15 percent ALND, 14 percent mastectomy, 26 percent lumpectomy) and all-cause mortality at 30 days (3.2 percent overall, 2 percent ALND, 4 percent mastectomy, 8.4 percent lumpectomy) and one year (31 percent overall, 29 percent ALND, 30 percent mastectomy, 41 percent lumpectomy).

"The highest mortality rate was associated with the least invasive procedure, lumpectomy, which appeared to be performed in the sickest patients," Tang said. "A higher mortality rate is somewhat expected due to advanced age and increased co-morbidities present in nursing home residents. However, a 30-day mortality of 8 percent is much higher than would be anticipated for a surgical procedure that is generally considered very low risk."

The researchers recommend that long-term nursing home residents with breast cancer consider hormonal therapy, such as endocrine therapy or radiotherapy, or symptom management only instead of surgery. Further studies should specifically evaluate this population in the outpatient setting and compare outcomes in those with and without the surgical intervention. Tools to support informed decision-making also are needed, they said.

The deferring of [breast cancer surgery](#) in frail elderly women is similar to prostate specific antigen (PSA) testing, Tang said. PSA testing is common in elderly men despite evidence that those without aggressive prostate cancer are unlikely to benefit from diagnosis and treatment and

may face significant risks for quality of life if they undergo prostate surgery.

"While some clinicians, patients and caregivers believe breast surgery is necessary to prevent morbidity and mortality from [breast cancer](#), the risks of harm may outweigh the benefit in this frail, vulnerable population, in which many have a limited life expectancy," said senior author Emily Finlayson, MD, MS, professor of surgery, health policy and geriatrics at UCSF.

Provided by University of California, San Francisco

Citation: Breast cancer surgery in frail elderly women linked to poor results (2018, August 29) retrieved 30 April 2024 from

<https://medicalxpress.com/news/2018-08-breast-cancer-surgery-frail-elderly.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--