

Cannabinoid in breast milk up to six days after marijuana use

August 28 2018



(HealthDay)—Most breast milk samples have measurable

Δ -9-tetrahydrocannabinol (Δ 9-THC) up to about six days after maternal use, according to a study published online Aug. 27 in *Pediatrics*.

Kerri A. Bertrand, M.P.H., from the University of California in San Diego, and colleagues quantified cannabinoids in [human milk](#) after maternal marijuana use. Fifty breastfeeding women who reported marijuana use provided 54 breast [milk](#) samples between 2014 and 2017.

The researchers found that up to about six days after last reported use, Δ 9-THC was detectable in 63 percent of the 54 samples; the median concentration was 9.47 ng/mL. Detectable levels of 11-hydroxy- Δ -9-tetrahydrocannabinol or cannabidiol were seen in five samples. No measurable Δ 9-THC was seen in the sample with the highest [concentration](#) of cannabidiol. A significant predictor of log Δ 9-THC concentrations was the number of hours since last use (-0.03). Other significant predictors of log Δ 9-THC concentrations were the number of daily uses and time from [sample collection](#) to analysis (0.51 and 0.08 , respectively), after adjustment for time since last use.

" Δ 9-THC was measurable in a majority of breast milk samples up to approximately six days after maternal marijuana use," the authors write.

More information: [Abstract/Full Text](#)
[Editorial](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Cannabinoid in breast milk up to six days after marijuana use (2018, August 28)
retrieved 25 April 2024 from
<https://medicalxpress.com/news/2018-08-cannabinoid-breast-days-marijuana.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.