

Does chemotherapy harm ability to function for older women with breast cancer?

August 28 2018



Credit: CC0 Public Domain

Older women are at higher risk for developing breast cancer than

younger women are—almost half of all breast cancer cases, and most breast cancer deaths, occur in women who are 65 or older. Despite this, we know very little about how breast cancer and its treatments affect older women. In particular, we don't fully understand how the disease and chemotherapy treatments affect a woman's ability to function and perform daily activities.

For older adults, knowing how chemotherapy may affect you is important, especially if there's a chance it could affect your ability to live independently. Understanding your risk for such problems would be good information to have when it comes to choosing treatments.

To learn more about how [breast cancer](#) and its treatments might affect [older women's](#) abilities to [function](#), a team of researchers designed a study. They published their results in the *Journal of the American Geriatrics Society*.

The researchers reviewed information from an earlier study, which included 633 women aged 65 or older who had been diagnosed with breast [cancer](#). That study compared the effectiveness of two different chemotherapy treatments. Researchers of that study asked participants questions, including:

- Do you have trouble with strenuous activities, like carrying a heavy shopping bag?
- Do you have any trouble taking a long walk?
- Do you have any trouble taking a short walk outside the house?
- Do you need to stay in bed or in a chair during the day?
- Do you need help with eating, dressing, washing yourself, or using the toilet?

In their new study, the researchers' goal was to learn about changes in the participants' ability to perform daily activities after the participants

received chemotherapy. The researchers also wanted to learn more about why some women lost some of their abilities—or experienced a decline—in performing daily activities, and why other women were able to recover the abilities they had before their [treatment](#).

The researchers said that their findings might help identify breast cancer survivors at risk of physical function decline and might shed light on future treatments designed to decrease this risk.

The researchers learned that short-term decline in physical function was common among older adults treated with chemotherapy for breast cancer. In the study, almost half (42 percent) of the participants experienced a decrease in their ability to function from pre-chemotherapy to the end of treatment. And almost one-third (30 percent) of the participants experienced a decline in their ability to function from before they started chemotherapy to 12 months later.

Among patients who experienced physical function decline from before chemotherapy to end of chemotherapy, about half (47 percent) were resilient, meaning they "bounced back" to the activity level they previously had within 12 months after starting chemotherapy treatment.

The impact of treatment on physical function is an important consideration for older adults. Functional decline is associated with loss of independence and increased risks for hospitalization and nursing home placement.

Interestingly, some older adults who experience problems performing their [daily activities](#) after cancer treatment are able to bounce back. Also known as "resilience," this process of recovery is considered vital to successful aging. Resilient [older adults](#) may be more able to recover after a stressful experience like chemotherapy.

"In our study, about half of the patients who experienced functional decline were able to 'bounce back' to their former function, and we considered them to be physically resilient," said the researchers. "We also learned that half of the patients were resistant to decline, and maintained their functional status throughout treatment," they added.

Risk factors for [functional decline](#) one year after [chemotherapy](#) included:

- Fatigue
- Having breathing difficulties
- Being unmarried or lacking social support, such as assistance from family and friends
- Poor appetite, which could indicate poor nutrition and/or depression

The helpful actions that could increase ability to "bounce back" include:

Diet and exercise programs. One study showed that starting a diet and exercise program within 18 months of breast cancer diagnosis is linked to an improvement in the ability to function.

Behavior Interventions. In one study, 49 percent of older, overweight survivors of breast, colorectal, and prostate cancer who enrolled in a two-year exercise, diet, and weight-loss program did not experience a decline in their abilities to function. In fact, 57 percent recovered to their pre-treatment level.

More information: Arti Hurria et al, Functional Decline and Resilience in Older Women Receiving Adjuvant Chemotherapy for Breast Cancer, *Journal of the American Geriatrics Society* (2018). [DOI: 10.1111/jgs.15493](https://doi.org/10.1111/jgs.15493)

Provided by American Geriatrics Society

Citation: Does chemotherapy harm ability to function for older women with breast cancer?
(2018, August 28) retrieved 20 April 2024 from

<https://medicalxpress.com/news/2018-08-chemotherapy-ability-function-older-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.