

Sleep disorder linked with abnormal lipid levels

August 22 2018

In the *Respirology* study, investigators identified strong associations between several measures of OSA severity and higher total cholesterol, higher LDL-cholesterol, elevated triglycerides, and lower HDL-cholesterol. Lipid status was influenced by geographical location with the highest total cholesterol concentration recorded in Northern Europe.

The analysis included 8592 adults across Europe who were not diagnosed with hyperlipidaemia and were not taking lipid-lowering drugs.

"Our data clearly suggest that sleep apnoea may have a negative impact on lipid levels, which may in part explain the association between sleep apnoea and increased risk for cardiovascular disease," said senior author Dr. Ludger Grote, of Gothenburg University, in Sweden. "Patients with [sleep apnoea](#) therefore need careful management of all [cardiovascular risk factors](#) including hyperlipidaemia."

More information: *Respirology* (2018).
onlinelibrary.wiley.com/doi/10.1111/resp.13372

Provided by Wiley

Citation: Sleep disorder linked with abnormal lipid levels (2018, August 22) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-08-disorder-linked-abnormal-lipid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.