


Drugs for heart failure are still under-prescribed, years after initial study

August 1 2018



Credit: CC0 Public Domain

A UCLA-led study found that many people with heart failure do not receive the medications recommended for them under guidelines set by the American College of Cardiology, American Heart Association and Heart Failure Society of America.

The research also found that doctors frequently prescribe medications at

doses lower than those recommended by the guidelines, especially for older people, those with kidney disease, those with worsening symptoms or those who were recently hospitalized for [heart failure](#). Further study is needed to determine why people in those four groups specifically were prescribed lower-than-recommended doses.

The study, which looked at the three categories of [heart](#) failure medications, found that between 27 percent and 67 percent of patients were not prescribed the recommended drugs. And when patients did receive the medications, they were generally at a lower-than-recommended dose. Less than 25 percent of patients simultaneously received all three medication types, and only 1 percent received the target doses of all three medication types.

About 5.7 million people in the United States have heart failure, according to a 2016 report by the American Heart Association. Heart failure is associated with a lower quality of life and frequent hospitalizations, and it contributes to more than 300,000 deaths each year in the U.S. In half of people with heart failure, the disease is caused by a weak heart muscle that prevents the heart from ejecting a normal amount of blood with each heartbeat, a condition called reduced ejection fraction.

Several medications have been proven in large clinical trials to help people with heart failure and reduced ejection fraction live longer and feel better. Research conducted between 2007 and 2009 showed that many patients were not receiving the recommended doses of these medications. The new study sought to determine if there have been improvements in prescribing practice as well as which patients are most likely to receive less [medication](#) than recommended.

The study included 3,518 [patients](#) from 150 primary care and cardiology practices who were enrolled in the Change the Management of Patients

with Heart Failure registry, or CHAMP-HF, a study of adult outpatients who were diagnosed with heart failure with reduced ejection fraction.

The results suggest that use and dosing of heart failure medications has not improved over the past decade. The report says new strategies are needed to more effectively achieve and maintain recommended doses of heart failure medications and that there is a substantial opportunity to improve dosing of heart failure medications, which would improve the care and outcomes for people with heart [failure](#).

The study appears in the July 24 issue of the *Journal of the American College of Cardiology*.

More information: Stephen J. Greene et al, Medical Therapy for Heart Failure With Reduced Ejection Fraction, *Journal of the American College of Cardiology* (2018). [DOI: 10.1016/j.jacc.2018.04.070](https://doi.org/10.1016/j.jacc.2018.04.070)

Provided by University of California, Los Angeles

Citation: Drugs for heart failure are still under-prescribed, years after initial study (2018, August 1) retrieved 2 May 2024 from

<https://medicalxpress.com/news/2018-08-drugs-heart-failure-under-prescribed-years.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--