


High-speed atomic force microscopy reveals clock protein interactions

August 21 2018, by Heidi Hall


Real-time Interaction of KaiA with native KaiC (CII)

Native KaiC
> 80% de-phosphorylated:



+ KaiA @ ~ 60 s
longer dwell time
(higher affinity)

Native KaiC
> 80% phosphorylated:



+ KaiA @ time 0
shorter dwell time
(lower affinity)

Johnson's team used high-speed atomic microscopy to see, in real time, how clock proteins interact. Credit: Carl Johnson Laboratory/Vanderbilt University

For the first time, researchers have seen how proteins involved in the daily biological clock interact with each other, helping them to further understand a process tied to numerous metabolic and eating disorders, problems with shift work, jet lag and mental health issues.


Working with the [clock proteins](#) KaiA, KaiB and KaiC from a model system, Carl Johnson, professor of biological sciences, and his team used high-speed [atomic force microscopy](#) to visualize the kinetics of the proteins' interactions. In the presence of adenosine triphosphate—the energy currency of living cells—the proteins rapidly and rhythmically bound to each other, leading to a regulative process called phosphorylation.

Their results, detailed in the article "Revealing circadian mechanisms of integration and resilience by visualizing clock proteins working in real time," were published Aug. 14 in the journal *Nature Communications*.

They discovered the on-and-off interactions between KaiA and KaiC take only seconds but combine to create a 24-hour oscillation of phosphorylation in a test tube. This biochemical clockwork keeps the rhythm going over at least 10 days without diminishing.

"We also learned through mathematical modeling that these interactions make the whole system more robust, no matter its environment," Johnson said. "Perhaps the environment inside cells is changing, but the clock can't keep accurate time if it changes with the intracellular environment. The high speed interactions between KaiA and KaiC that we observed keeps the oscillation resilient and accurate."

He said using clock proteins from model organisms allows more rapid progress because the system is relatively simple and therefore easier to see the key interactions.



With on-and-off interactions with other proteins that take only seconds, KaiA constitutes its own internal clock and keeps the rhythm going over at least 10 days without losing strength. Credit: Carl Johnson Laboratory/Vanderbilt University

More information: Tetsuya Mori et al. Revealing circadian mechanisms of integration and resilience by visualizing clock proteins working in real time, *Nature Communications* (2018). [DOI: 10.1038/s41467-018-05438-4](https://doi.org/10.1038/s41467-018-05438-4)

Provided by Vanderbilt University

Citation: High-speed atomic force microscopy reveals clock protein interactions (2018, August 21) retrieved 27 April 2024 from <https://medicalxpress.com/news/2018-08-high-speed-atomic-microscopy-reveals-clock.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.