

Sleeping five hours or less a night associated with doubled risk of cardiovascular disease

August 26 2018



Credit: CC0 Public Domain

Middle-aged men who sleep five hours or less per night have twice the risk of developing a major cardiovascular event during the following two decades than men who sleep seven to eight hours, according to research

presented today at ESC Congress 2018.

Study author Ms Moa Bengtsson, of the University of Gothenburg, Sweden, said: "For people with busy lives, sleeping may feel like a waste of time but our study suggests that [short sleep](#) could be linked with future [cardiovascular disease](#)."

Previous studies have generated conflicting evidence on whether short sleep is associated with a greater chance of having a future cardiovascular event. This study investigated this relationship in 50-year-old men.

In 1993, 50% of all men born in 1943 and living in Gothenburg were randomly selected to participate in the study. Of the 1,463 invited, 798 (55%) men agreed to take part. Participants underwent a physical examination and completed a questionnaire on current health conditions, average sleep [duration](#), physical activity, and smoking. The men were divided into four groups according to their self-estimated average sleep duration at the start of the study: five or less hours, six hours, seven to eight hours (considered normal sleep duration), and more than eight hours.

Participants were followed-up for 21 years for the occurrence of major [cardiovascular events](#), which included heart attack, stroke, hospitalisation due to heart failure, coronary revascularisation, or death from cardiovascular disease. Data on cardiovascular events were collected from medical records, the Swedish Hospital Discharge Registry, and the Swedish Cause of Death Register.

Men with incomplete data on sleep duration, incomplete follow-up information, or who had a major cardiovascular event before the start of the study were excluded, leaving a total of 759 men for the analyses.

High blood pressure, diabetes, obesity, current smoking, low physical activity, and poor sleep quality were more common in men who slept five or fewer hours per night compared to those who got seven to eight hours.

Compared to those with normal sleep duration, men who slept five or fewer hours per night had a two-fold higher risk of having a major cardiovascular event by age 71. The risk remained doubled after adjusting for [cardiovascular risk factors](#) at the start of the study including obesity, diabetes, and smoking.

Ms Bengtsson said: "Men with the shortest [sleep duration](#) at the age of 50 were twice as likely to have had a cardiovascular event by age 71 than those who slept a normal amount, even when other risk factors were taken into account."

She continued: "In our study, the magnitude of increased cardiovascular risk associated with insufficient sleep is similar to that of smoking or having diabetes at age 50. This was an observational study so based on our findings we cannot conclude that short sleep causes cardiovascular disease, or say definitively that sleeping more will reduce risk. However, the findings do suggest that sleep is important—and that should be a wake-up call to all of us."

More information: Abstract: "Middle age men with short sleep duration have two times higher risk of cardiovascular events than those with normal sleep duration, a cohort study with 21 years follow-up"

Provided by European Society of Cardiology

Citation: Sleeping five hours or less a night associated with doubled risk of cardiovascular

disease (2018, August 26) retrieved 27 April 2024 from
<https://medicalxpress.com/news/2018-08-hours-night-cardiovascular-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.