

Novel imaging biomarker to help predict coronary inflammation

August 28 2018



Credit: CC0 Public Domain

Researchers at Cleveland Clinic, University of Oxford and University of Erlangen have identified a novel imaging biomarker, which has been found to be able to predict all-cause and cardiac mortality by measuring inflammation of fatty tissue surrounding the coronary arteries.

Coronary artery inflammation inhibits [fatty tissue](#) formation surrounding the blood vessels, known as perivascular fat. Researchers developed the perivascular fat attenuation index (FAI) as an imaging biomarker to quantify inflammation-induced changes in perivascular fat. FAI captures coronary inflammation by mapping the changes in perivascular fat on coronary computed tomography angiography (CTA), enabling early detection of coronary inflammation.

"This is an exciting new technology which has the potential for providing a simple, non-invasive answer to detect patients at risk for future fatal [heart](#) attacks, said co-first author Milind Desai, M.D., Cleveland Clinic cardiologist. "More importantly, it highlights the incredible value of cross-continent collaboration to validate the findings in different populations."

Cardiovascular Risk Prediction using Computed Tomography (CRISP-CT) study collected data from the two cohorts of consecutive patients undergoing coronary CT angiography—1,872 patients in Germany from 2005 to 2009 (derivation cohort) and 2,040 [patients](#) at Cleveland Clinic from 2008 to 2016 (validation cohort). Median patient age in the cohorts was 62 and 53 years.

In both cohorts, higher perivascular FAI values—indicating greater coronary [inflammation](#)—were associated with significantly higher rates of death from any cause and death from cardiac causes.

"This new technology may prove transformative for primary and secondary prevention. For the first time we have a set of biomarkers, derived from a routine test that is already used in everyday clinical practice, that measures what we call the 'residual cardiovascular risk', currently missed by all risk scores and non-invasive tests," said Charalambos Antoniades, M.D., who led the study at the University of Oxford's Division of Cardiovascular Medicine. "Knowing who is at

increased risk for a heart attack could allow us to intervene early enough to prevent it. I expect these biomarkers to become an essential part of standard CT coronary angiography reporting in the coming years."

According to the CDC, about 610,000 die of heart disease in the United States every year. Coronary heart disease (CHD) is the most common type of heart disease, and about 735,000 Americans suffer from heart attack each year.

The research was presented as a late-breaking presentation at the European Society of Cardiology Congress 2018 in Munich, Germany and simultaneously published online in *The Lancet*.

Provided by Cleveland Clinic

Citation: Novel imaging biomarker to help predict coronary inflammation (2018, August 28) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-08-imaging-biomarker-coronary-inflammation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.