

## The risk of lung cancer for young breast cancer survivors

August 7 2018, by Jean-Philippe Pignol And Nienke Hoekstra



Credit: AI-generated image (disclaimer)

Women who have been lucky enough to survive breast cancer may face increased risk of lung cancers, <u>according to a new study in *Radiotherapy*</u> <u>and Oncology</u>. The study shows how this can be prevented—limiting the radiation dose to a lung, using a treatment which targets only a small part of the breast.



Breast cancer is the most frequently diagnosed cancer in women, with more than 266,000 new cases diagnosed in 2018 in the United States.

Thanks to the widespread use of screening mammography, most women are now diagnosed at an early stage —meaning that the cancer has only grown in the breast and not spread under the armpit or to distant organs. These individuals have a good prognosis with <u>only a 1.3 per cent risk of dying of cancer recurrence after five years</u>.

Treatment currently involves a limited breast surgery followed by multiple daily <u>radiation</u> treatments targeting the whole breast. However, despite being cautiously delivered, a small amount of radiation can spill into the lung.

Using a complex predictive model called the <u>Biologic Effects of Ionizing</u> <u>Radiation (BEIR) VII</u>, and by testing multiple radiation techniques on a realistic model of a <u>breast cancer</u> patient, the study by Ph.D. student <u>Nienke Hoekstra</u> shows that the risk of <u>lung cancer</u> slowly increases five to 10 years after a breast radiation treatment. It reaches three per cent of surviving patients after more than 30 years.

## A small part of the breast

This risk of lung cancer is a real concern for young patients diagnosed with early stage breast cancers —for example, a ductal carcinoma in situ (DCIS), which is a form of non-invasive cancer and hence a very highly curable disease. These patients have the chance of living long enough to experience this scary treatment side-effect.

The good news is that risk is not the same for all radiotherapy techniques. The study shows that a new form of radiotherapy for earlystage breast cancers —called accelerated partial breast irradiation (APBI) —dramatically reduces this risk.



APBI delivers the radiation treatment to only a small portion of the breast. It focuses on the cavity left after the surgery, where there is risk that cancer cells may remain.



Robotic delivery of radiation from all angles around the breast is the focus of ongoing research at Erasmus University. Author provided

There are multiple advantages to APBI, including the delivery of treatment in just five days. And the study found it results in much lower radiation exposure to body organs, including the lungs, than conventional treatments.



APBI is a technique on the rise. This study adds to the growing amount of evidence that it should be the preferred technique for women with very low-risk breast cancer.

But not all patients will benefit from APBI. Some need more complex and extensive radiotherapy to save their lives. Patients should be encouraged to discuss this with their radiation oncologist.

## **Tiny radioactive seeds**

Among APBI techniques tested, <u>"brachytherapy"</u> —a form of radiotherapy where radioactive material is implanted inside the surgical cavity —appears to be the safest.

A particular form of brachytherapy developed in Canada, called "permanent breast seed implant," is the one with the lowest risk of secondary lung cancer.

This technique, which is also used to treat prostate cancer, <u>involves the</u> <u>permanent implantation of tiny radioactive seeds</u>, the size of a grain of rice, during a one-hour procedure performed under light sedation. Patients are discharged the same day.

The radiation emitted by the radioactive seeds is so weak that it is absorbed within a few millimetres and hence rarely reaches other organs.

The good news is that there is still lots of room for improvement —for example, using a robotic delivery of radiation from all angles around the breast to better avoid the lung. This is the focus of ongoing research.

This article was originally published on <u>The Conversation</u>. Read the <u>original article</u>.



## Provided by The Conversation

Citation: The risk of lung cancer for young breast cancer survivors (2018, August 7) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2018-08-lung-cancer-young-breast-</u> <u>survivors.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.