

# How meditation affects the brain to help you stress less

August 22 2018, by Michaela Pascoe

---



Credit: AI-generated image ([disclaimer](#))

In Australia, about [one in six adults](#) practise meditation, while [one in ten](#) practise yoga. People often turn to yoga or meditation as a way to take time out and manage the stress of their day-to-day lives.

Stress is common, and ongoing stress can contribute to the onset of [a](#)

[range of psychological issues](#), such as depression and anxiety.

[Meditation](#) and [yoga](#) have been shown to reduce people's self-reported levels of stress. This is likely due, at least in part, to the effects that meditation and [yoga](#) have on the brain's [stress response](#) system.

## **How the brain responds to stress**

The body's automatic stress response is controlled by the autonomic nervous system. The autonomic nervous system plays a key role in stress reactivity via its two main divisions: the sympathetic nervous system and the parasympathetic nervous system.

A main function of the the sympathetic nervous system is to mobilise the body to fight or flee from stressful or threatening situations, via control of internal muscles, organs and glands. This is called the "fight or flight" response.

The parasympathetic nervous system counterbalances the sympathetic nervous system and returns the body to its natural baseline state after the systematic nervous system activates.

In many cases the parasympathetic nervous system and sympathetic nervous system have opposing but complementary functions. For example, the sympathetic nervous system increases heart rate, [blood pressure](#) and the downstream release of stress-related hormones such as cortisol, whereas the parasympathetic nervous system decreases all of these factors.

So by measuring these [we can identify](#) if people are experiencing a homeostatic state or a more stressful state, on a physiological level.

We [reviewed](#) how yoga and different forms of meditation influence the

brain's stress response system by studying physiological markers of stress.

## What are the different forms of meditation?

A common method of classifying meditation techniques distinguishes between [open monitoring](#), [focused attention](#), and automatic [self-transcending](#) meditation.

Open monitoring or mindfulness-based meditations involve the practice of observing the content of our ongoing experience in a non-reactive way, [to become reflectively aware](#) of cognitive and emotional patterns. Instead of focusing attention on a particular object, the meditator aims to pay attention to and monitor all aspects of experience as they come up, without judgement or attachment. An example would be feeling the sensation of the seat beneath you while meditating.

In focused attention meditation, attention is focused and sustained [on a particular object](#) and brought back to the object when the mind has wandered. In this way, the meditator is controlling their own attention. The object the person focuses on may be the breath, a mantra, visualisation, a part of the body, or an external object. Each time the meditator notices that their attention wanders, they actively bring it back to their object of attention.

Automatic self-transcending involves the use of a mantra, usually Sanskrit sounds, which the meditator can attend to without effort or concentration. The aim is that the mantra becomes secondary and ultimately disappears as self-awareness increases. In automatic self-transcending meditation, the mind should be [free from focus](#) and mental effort. It is practised for 15–20 minutes twice a day while sitting with closed eyes.

## What the evidence says

We found that meditation and yoga reduce [diastolic blood pressure](#) (the lower range) by 3-8 millimetres of mercury (mmHg), compared with people who engaged in another activity, such as aerobic exercise or relaxation.

Both focused attention and automatic self-transcending meditation styles, as well as yoga, reduced systolic blood pressure (the upper range) by 4-5mmHg, compared with people who were not practising any kind of meditation or yoga. This is important because reductions in systolic and diastolic blood pressure of as little as two mmHg can [reduce the incidence](#) of heart disease and stroke.

Open monitoring and focused attention meditation and yoga reduced [heart rate](#) by three to four beats per minute. This is similar to the effects of aerobic exercise, which [reduced heart rate](#) by five beats per minute in one study.

Focused attention meditations and yoga both decreased measures of cortisol.

Our findings indicate that all forms of meditation studied reduce physiological stress markers in one way or another, and therefore, all forms are likely beneficial in managing stress.

In terms of deciding what form is best for reducing [stress](#), we would suggest practising a form that is enjoyable and therefore you will practise regularly and in an ongoing manner.

While understanding the different types of meditation is useful, meditation classifications [should not be considered](#) to be mutually exclusive, either within a single [meditation](#) session or over a lifetime of

practice. Most meditative techniques lie [somewhere on a continuum](#) between open monitoring and focused [attention](#) types.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: How meditation affects the brain to help you stress less (2018, August 22) retrieved 3 May 2024 from <https://medicalxpress.com/news/2018-08-meditation-affects-brain-stress.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.