

## **Training for parents referred to CPS improves toddler's physiological regulation**

August 28 2018, by Andy Fell

A parental training program for families referred to Child Protective Services improved toddlers' unconscious reactions to mildly stressful situations, as well as improving parents' behavior, according to a new study by researchers at the University of California, Davis and the University of Washington. The work is published August 28 in *Developmental Science*.

"In everything we do, the brain and body are coordinated through the autonomic nervous system," said Paul Hastings, professor in the UC Davis Department of Psychology and Center for Mind and Brain. "We know that these neurobiological systems are shaped by our experiences in the world, including parenting."

The autonomic nervous system functions primarily below the level of conscious thought and has two arms: the sympathetic and parasympathetic systems. The better-known sympathetic system regulates "fight or flight" responses. The parasympathetic system tends to calm those responses and contributes to our ability to engage socially and emotionally with other people.

People who have experienced maltreatment as <u>children</u> are at risk of developing depression and other <u>mental health problems</u> later in life. This may be tied to the effects of poor parenting on physiology, including the parasympathetic system. But under normal circumstances, it's difficult to study whether parenting style can cause changes in a child's physiology—you couldn't ethically or practically do an



experiment in which children are assigned to be in families with better or worse <u>parents</u> and see what happens, Hastings said.

## **Intervention to improve parenting**

Promoting First Relationships is a <u>training</u> program designed to improve sensitivity towards infants and toddlers in troubled families. Professor Monica Oxford of the University of Washington led a study of the program's effectiveness with 247 families who had been referred to Child Protective Services for maltreatment. Families either took the Promoting First Relationships training, which involves in-home visits and parent training sessions, or the standard service of receiving referrals and resources. That study showed that the training did improve parents' understanding and sensitive engagement with their toddlers and lowered the number of foster care placements.

Hastings' team used a sample of 29 families from the Promoting First Relationships training and 30 families from the comparison group to see if the treatment program that changed parenting behavior also had an effect on children's parasympathetic regulation, measured based on changes in heart rate in response to mildly challenging or stressful tasks such as completing a simple puzzle or petting a plastic spider that jumps when triggered. Typically, children react to such situations with a mild decrease in parasympathetic regulation, becoming more alert and ready to take in information. The team assessed the children at the end of the Promoting First Relationships trial, and again three and six months later.

Children of parents who received the training responded to the tasks with the typical pattern of mild parasympathetic decreases that indicated they were attentive to and engaged with the tasks. This response was strongest in the children of parents who showed the most improvement in sensitive behavior after the training.



Children of parents in the comparison group had a bigger drop in parasympathetic activity, which would open the door for their sympathetic system to drive fight-or-flight responses like becoming distressed and withdrawing. This response matches what earlier studies had found in adolescents and adults who had experienced maltreatment during their childhoods.

## **Inner strength**

Showing that an intervention that improves parenting behavior also improves young children's physiological regulation is important, Hastings said, because that capacity for regulation is an internal strength that children can take forward into their future experiences, which should help them to be resilient and to thrive.

This study worked with a particularly at-risk population, but helping parents who are within the normal range of behavior to become more sensitive and responsive could be beneficial for all children, Hastings said.

"Parents are heavy consumers of self-help books and websites, but we don't know how beneficial these things are for children's overall development," he said. "We know that parenting affects neurobiology and predicts later social and emotional adjustment, but we need evidence that programs like this make a difference in order to give parents the right advice."

Provided by UC Davis

Citation: Training for parents referred to CPS improves toddler's physiological regulation (2018, August 28) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2018-08-parents-cps-toddler-physiological.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.