

Low plasma levels of omega-3 fatty acids associated with preterm birth

August 3 2018



Credit: CC0 Public Domain

Pregnant women who had low plasma levels of long chain n-3 fatty acids in their first and second trimesters were at a significantly higher risk of early preterm birth when compared with women who had higher levels

of these fatty acids, according to new research from Harvard T.H. Chan School of Public Health in collaboration with Statens Serum Institut in Copenhagen. The study suggests that low concentrations of certain long chain fatty acids—eicosapentaenoic acid and docosahexaenoic acid (EPA+DHA)—may be a strong risk factor for preterm birth.

"At a time when many pregnant women are hearing messages encouraging them to avoid intake of fish altogether due to mercury content, our results support the importance of ensuring adequate intake of long chain omega-3 [fatty acids](#) in pregnancy. Consumers should consult the guidance issued last year by the U.S. Food and Drug Administration and Environmental Protection Agency to make informed choices about the best types of fish to consume and avoid in pregnancy," said lead author Sjurður F. Olsen, adjunct professor of nutrition at Harvard Chan School and head of the Centre for Fetal Programming at Statens Serum Institut in Copenhagen, Denmark.

The study will be published online August 3, 2018, in *EBioMedicine*.

Preterm birth is a leading cause of neonatal death and is associated with cognitive deficiencies and cardiometabolic problems later in life among survivors. For decades, it has been hypothesized that high intake of EPA+DHA, which is found in cold-water fish such as Atlantic mackerel, anchovies, salmon, and tuna and also in leaner species such as cod and haddock, can reduce the risk of preterm birth. While some studies have supported this hypothesis, research findings have been inconsistent.

For this new study, researchers examined data from the Danish National Birth Cohort, a nationwide study that follows 96,000 children in Denmark through questionnaires and registry linkages. They analyzed blood samples from 376 women who gave birth very prematurely (prior to 34 weeks of gestation) between 1996 and 2003 and 348 women who had a full-term birth. All of the women gave blood samples during their

first and second trimesters of pregnancy.

Analysis of the blood samples showed that women who were in the lowest quintile of EPA+DHA serum levels—with EPA+DHA levels of 1.6% or less of total plasma fatty acids—had a 10 times higher risk of early preterm birth when compared with women in the three highest quintiles, whose EPA+DHA levels were 1.8% or higher. Women in the second lowest quintile had a 2.7 times higher risk compared with women in the three highest quintiles.

The findings suggest that, among pregnant women with low levels of EPA+DHA, eating more fish or taking a fish oil supplement could potentially lower the risk of preterm birth. The authors cautioned, however, that broad generalizations about the study's findings may be limited due to the fact that it was conducted in Denmark, where preterm birth rates are low, and said the results should be replicated in other populations. They also cautioned that the findings may not solely reflect a variation in diet; variation in underlying genetic factors may also play a role.

"An effect of this magnitude is rare, but the precision of the estimate is tight, which supports the reliability these findings. It will be important to replicate these findings in other populations, but the results of this study certainly suggest that assessment of plasma EPA+DHA status in women has the potential to be used in the future to help predict women's risk," said co-author Jeremy Furtado, senior research scientist at Harvard Chan School.

"Early preterm birth has immense health, economic, and emotional costs. Our findings are consistent with the results of most randomized trials of long chain omega-3 fatty acid supplements in pregnancy and support the importance of ensuring adequate intake of these nutrients during pregnancy, either through fish intake or supplements, to help

prevent early [preterm birth](#)," said co-author Andrew Thorne-Lyman, an associate research scientist at Johns Hopkins Bloomberg School of Public Health who worked on this study while a faculty member at Harvard Chan School.

More information: *EBioMedicine* (2018). [DOI: 10.1016/j.ebiom.2018.07.009](#)

Provided by Harvard T.H. Chan School of Public Health

Citation: Low plasma levels of omega-3 fatty acids associated with preterm birth (2018, August 3) retrieved 3 May 2024 from <https://medicalxpress.com/news/2018-08-plasma-omega-fatty-acids-preterm.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--