

Predicting work status in patients with schizophrenia

August 21 2018




Figure 1: Analysis steps. Credit: Sumiyoshi, C., et al., Schizophr. Res. (2018)

Osaka University-led Japanese researchers clarified that intellectual deterioration (IQ decline) was related to work status in patients with

schizophrenia. The researchers also proposed a method for estimating probabilities of work outcome in those patients based on related factors, such as IQ decline, social function, and psychiatric symptoms.

Japanese scientists demonstrate that the difference between current full scale IQ and estimated premorbid IQ is related to work outcome in patients with [schizophrenia](#)

Schizophrenia is a mental disorder that occurs in one out of every 100 people. Its symptoms include positive symptoms, such as hallucinations and delusions, and negative symptoms, such as social withdrawal, apathy, and cognitive impairments. In many cases, these symptoms become chronic and relapse. In addition, many patients with schizophrenia show deterioration of intelligence, which prevents them from independent living in their community, and joining the work force, in particular.

A research team led by Ryota Hashimoto at Osaka University clarified that IQ decline, defined as a difference between current full scale IQ (FIQ) and estimated premorbid IQ, was related to work status in patients with schizophrenia. The researchers also proposed a method for estimating probabilities of work outcome in those patients based on related factors, such as IQ decline, social function, and [psychiatric symptoms](#). Their research results were published in Schizophrenia Research.

Predictive factors:

Intelligence decline
Social function
Psychiatric symptoms



Work outcomes:

Capacity for working longer than a certain amount of time dichotomized by criteria (0, 10, 20, or 30 hours per week)

Figure 2: Significant factors for predicting work outcomes. Credit: Fukushima University

First, various variables were compared between the preserved group (patients with preserved IQ), the deteriorated group (patients whose current intelligence was lower than their estimated premorbid intelligence), and the healthy controls (the group of healthy volunteers).

Next, they performed [logistic regression](#) analyses to predict work status in patients with schizophrenia using the significant variables found in the group comparisons. The work status was dichotomized by a criterion of 1, 10, 20, or 30 hours per week. At each criterion, patients were classified into either the above or the below criterion according to their actual work hours.

The analyses demonstrated that IQ decline was effective for predicting work status. Based on the equations obtained from the logistic regression models, they also presented a method for predicting probabilities for working longer than each criterion.

- An example from the Preserved group
 - Adjusted SFS 156
 - IQ decline 1.68
 - PANSS Total 61
 - Estimated probability=0.96

- An example from the Deteriorated group
 - Adjusted SFS 139
 - IQ decline -27.7
 - PANSS Total 83
 - Estimated probability=0.53




Figure 3: Examples of estimation for the less than 20 vs. 20 or longer (hours/week) criterion. Credit: Sumiyoshi, C., et al., *Schizophr. Res.* (2018)

The first author Chika Sumiyoshi at Fukushima University says, "The amount of work performed by patients with schizophrenia was predicted by doctors only based on their experience and intuition. Our [research results](#) will help them to explain how many hours patients can work to them and their families based on objective data and promote consensus-building between the patient and physician. This will help motivate patients to receive treatment and promote their reintegration into society.

More information: Chika Sumiyoshi et al. Predicting work outcome in patients with schizophrenia: Influence of IQ decline, *Schizophrenia Research* (2018). [DOI: 10.1016/j.schres.2018.05.042](https://doi.org/10.1016/j.schres.2018.05.042)

Provided by Osaka University

Citation: Predicting work status in patients with schizophrenia (2018, August 21) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-08-status-patients-schizophrenia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.