

## Stress gene influences chronic pain after car crash

August 27 2018



Credit: Linnstaedt et al., JNeurosci (2018)



A study of more than 1,000 motor vehicle accident survivors published in *JNeurosci* reveals a common variant in a gene involved in the stress response that increases vulnerability to developing chronic pain. Addressing the interaction between this genetic variant and posttraumatic stress may represent a supplemental or alternative to treatment with addictive opioids.

Sarah Linnstaedt and colleagues conducted two prospective studies of European American and African American men and women admitted to the emergency room following a <u>motor vehicle</u> collision.

In addition to genotyping these patients, the researchers assessed their distress immediately after the accident as well as their pain and post-traumatic stress symptoms six weeks later.

Participants of both ethnicities with a particular variant in the gene FKBP5 reported more severe pain at follow up, which was related to their level of distress. Further analyses uncovered a molecular mechanism by which this variant increases FKBP5 expression and mediates its relationship with post-traumatic chronic pain.

**More information:** *JNeurosci* (2018). <u>DOI:</u> <u>10.1523/JNEUROSCI.3458-17.2018</u>

Provided by Society for Neuroscience

Citation: Stress gene influences chronic pain after car crash (2018, August 27) retrieved 7 May 2024 from <u>https://medicalxpress.com/news/2018-08-stress-gene-chronic-pain-car.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.