


Mixed trends in teenage 'new smoker' rates in Europe

August 22 2018



Crude rates of smoking initiation according to age, by sex and by decade. Credit: Marcon et al., 2018

In most of Europe, the rates of smoking initiation among older teens

have declined since the 1970s, while "new smoker" rates among younger teens have risen in recent years, according to a study published August 22, 2018 in the open-access journal *PLOS ONE* by an international team of researchers involved in the Ageing Lungs in European Cohorts (ALEC) study, coordinated by Deborah Jarvis, Imperial College of London, UK.

Smoking is the leading cause of avoidable mortality and the strongest modifiable risk factor for a number of diseases. Many smokers initially become addicted to nicotine during adolescence, making it critical to understand [smoking](#) trends in this population.

The ALEC study pooled data from six multicenter studies, including 119,104 people from 17 countries. Information on smoking was retrospectively obtained from participants, who answered questions including "Are you a smoker," "Have you ever smoked for as long as a year?" and "How old were you when you started smoking?" Data was grouped by region of Europe and analyzed both as a whole and by [region](#)

Overall, the rates of smoking initiation before 1970 peaked at age 18 in males and 19 in females; in the 2000s they peaked at ages 16 and 15. For both sexes, and all regions, smoking initiation during late adolescence (age 16 to 20) declined from the 1970s through to the 2000s. In the late 2000s, rates of smoking initiation in late adolescence were lowest in North Europe, with 20 people per 1,000 per year initiating smoking. In other regions, rates remained high at 40-80 new smokers per 1000 per year. Smoking initiation rates during [early adolescence](#) (age 11 to 15) increased in nearly all regions from 1990 through the 2000s. In West Europe, smoking initiation rates in this population reached 40 people per 1,000 per year around 2005.

"Smoking initiation is still unacceptably high among European

adolescents, and increasing rates among those aged 15 or less deserve attention," the authors say. "Reducing initiation in [adolescents](#) is fundamental, since youngsters are particularly vulnerable to nicotine addiction and tobacco adverse effects."

First author Alessandro Marcon adds: "Unacceptably high rates of [smoking initiation](#) during late adolescence by the late 2000s, and increasing rates in children from 1990 on, suggest the necessity for most European countries to intensify the implementation of tobacco control measures reaching the youngsters."

More information: Marcon A, Pesce G, Calciano L, Bellisario V, Dharmage SC, Garcia-Aymerich J, et al. (2018) Trends in smoking initiation in Europe over 40 years: A retrospective cohort study. *PLoS ONE* 13(8): e0201881. doi.org/10.1371/journal.pone.0201881

Provided by Public Library of Science

Citation: Mixed trends in teenage 'new smoker' rates in Europe (2018, August 22) retrieved 24 April 2024 from <https://medicalxpress.com/news/2018-08-trends-teenage-smoker-europe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--