

New blood test detects early stage pancreatic cancer

September 17 2018



Axial CT image with i.v. contrast. Macrocystic adenocarcinoma of the pancreatic head. Credit: public domain

Pancreatic cancer is currently very difficult to detect while it is still resectable. A new blood test developed by researchers at Lund University in Sweden, Herlev Hospital, Knight Cancer Center and



Immunovia AB, can detect pancreatic cancer in the very earliest stages of the disease. The results have been published in the *Journal of Clinical Oncology*.

Due to diffuse symptoms, <u>pancreatic cancer</u> is usually diagnosed very late in the disease progression. Therefore, despite pancreatic cancer representing less than 3% of all cancer cases, more people currently die from it than breast cancer. By 2030, pancreatic cancer is expected to be the second deadliest type of cancer in the world.

"Our test can detect pancreatic cancer with 96% accuracy at <u>stage</u> I and II, while there is still the possibility of successful surgical intervention. There is currently no cure and few treatment options for advanced pancreatic cancer, which is the late stage when pancreatic cancer is usually diagnosed", explains Carl Borrebaeck, professor at the department of Immunotechnology at Lund University.

The study used samples from patients in both Denmark and the US, at different stages of the disease.

The <u>blood test</u> is developed on a so-called antibody microarray that consists of hundreds of recombinant antibody fragments. These antibody fragments are specific for a number of immune-regulatory proteins, cancer-associated antigens, and so on.

Since the immune system is the first to respond to threats like complex diseases, such as cancer, autoimmune diseases and infections, the microarray was designed to mirror this early response. This provides information about the development of tumours long before being visible on CT or detected by ctDNA. From those hundreds of markers, 29 markers were selected to detect pancreatic cancer with 96% accuracy at stage I and II.



In the future, the screening method could be used to screen people who are at a higher risk of developing <u>pancreatic cancer</u>, such as those with a hereditary risk, newly onset diabetes patients and patients with chronic inflammation of the pancreas.

The next step has already been initiated, which is a large US prospective study for high risk individuals.

More information: Linda D. Mellby et al. Serum Biomarker Signature-Based Liquid Biopsy for Diagnosis of Early-Stage Pancreatic Cancer, *Journal of Clinical Oncology* (2018). DOI: 10.1200/JCO.2017.77.6658

Provided by Lund University

Citation: New blood test detects early stage pancreatic cancer (2018, September 17) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2018-09-blood-early-stage-pancreatic-</u> <u>cancer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.