


Research suggests increased cortical activation in cannabis users' brains in resting state

September 5 2018



Left hemisphere of J. Piłsudski's brain, lateral view. Credit: public domain

Recent research from the Center for BrainHealth at The University of Texas at Dallas shows that cannabis users experience increased cortical activation during the brain's resting state when compared with nonusers.

The resulting "noisy [brain](#)" might impair brain activity and disrupt cognitive processes, said Dr. Shikha Prashad, the study's lead author and a research scientist at the Center for BrainHealth.

"This study is the first to characterize global [cortical activation](#) and both inter- and intrahemispheric functional connectivity during resting state in [cannabis users](#)," said Prashad, who works in the lab of Dr. Francesca M. Filbey, the Bert Moore Chair in BrainHealth at UT Dallas.

The results of the study were published in the journal *NeuroImage*.

The researchers collected electroencephalogram (EEG) data from 38 participants in two groups: 17 cannabis users and 21 nonusers. The EEG test measures electrical activity in the brain—also known as brain waves. Scientists measured the synchronization of brain waves to evaluate the strength of the brain signals in different cortical regions. Such coherence of the EEG signal is a measure of the communication between different regions of the brain.

The results demonstrated that cannabis users exhibited increased synchronization—or activation—of most of the different types of [brain waves](#), compared to nonusers. Similar results have been found in other studies of heroin-, alcohol- and cocaine-dependent users.

In addition, greater cortical communication among the frontal regions of the brain was observed in cannabis users. Prashad said this could signify that participants had difficulty inhibiting neural activity that has been observed in Dr. Filbey's previous studies, which would cause them to exert more effort as they attempted to stop doing certain tasks. Changes in communication between the brain's [cortical areas](#) could also be related to cognitive impairments correlated with cannabis use.

The findings of this research suggest that further studies are needed

during task-related activity for comparison, Prashad said. These results could contribute to understanding cognitive impairments and the development of electrophysiological patterns to help map and evaluate the success of interventions.

More information: Shikha Prashad et al, Cannabis users exhibit increased cortical activation during resting state compared to non-users, *NeuroImage* (2018). [DOI: 10.1016/j.neuroimage.2018.06.031](https://doi.org/10.1016/j.neuroimage.2018.06.031)

Provided by Center for BrainHealth

Citation: Research suggests increased cortical activation in cannabis users' brains in resting state (2018, September 5) retrieved 28 April 2024 from <https://medicalxpress.com/news/2018-09-cortical-cannabis-users-brains-resting.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.