

Crunched for time? High-intensity exercise = same cell benefits in fewer minutes

September 20 2018



Credit: CC0 Public Domain

A few minutes of high-intensity interval or sprinting exercise may be as effective as much longer exercise sessions in spurring beneficial improvements in mitochondrial function, according to new research. The

small study is published ahead of print in the *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology*.

Mitochondria, the energy centers of the cells, are essential for good health. Previous research has found that exercise creates new mitochondria and improves the function of existing mitochondria. Altered [mitochondrial function](#) in response to a single session of exercise generates signals that may lead to beneficial changes in the cells, lowering the risk for chronic disease. High-intensity interval exercise consists of short bursts of high-intensity aerobic exercise—physical activity that raises the heart rate—alternating with brief recovery periods. Whether the intensity of a workout affects [mitochondrial response](#) is unclear.

A team of researchers studied eight young adult volunteers as they participated in cycling workouts of varying intensity.

- Moderate intensity consisted of 30 minutes of continuous exercise at 50 percent peak effort.
- High-intensity interval exercise consisted of five four-minute cycling sessions at 75 percent peak effort, each separated by one minute of rest.
- Sprint cycling consisted of four 30-second sessions at maximum effort, each separated by 4.5 minutes of recovery time.

The research team measured the amount of energy the volunteers spent on each workout and compared mitochondrial changes in the participants' thigh muscles before and after each [exercise session](#). The researchers found that levels of hydrogen peroxide—a type of molecule involved in cell signaling called "reactive oxygen species" that contains oxygen and hydrogen—in different parts of the mitochondria change after exercise. While too much [reactive oxygen species](#) can be damaging to the [cells](#), the researchers noted that the volunteers' levels were an

appropriate amount to potentially promote cell responses that benefit metabolic function rather than cause damage.

In addition, the research team found that fewer minutes of higher-intensity exercise produced similar mitochondrial responses compared to a longer moderate-intensity activity. "A total of only two minutes of sprint interval exercise was sufficient to elicit similar responses as 30 minutes of continuous moderate-intensity aerobic exercise," the researchers wrote. "This suggests that exercise may be prescribed according to individual preferences while still generating similar signals known to confer beneficial metabolic adaptations. These findings have important implications for improving our understanding of how [exercise](#) can be used to enhance metabolic health in the general population."

More information: Adam James Trewin et al, Acute HIIE elicits similar changes in human skeletal muscle mitochondrial H₂O₂ release, respiration and cell signaling as endurance exercise even with less work, *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* (2018). [DOI: 10.1152/ajpregu.00096.2018](https://doi.org/10.1152/ajpregu.00096.2018)

Provided by American Physiological Society

Citation: Crunched for time? High-intensity exercise = same cell benefits in fewer minutes (2018, September 20) retrieved 30 April 2024 from <https://medicalxpress.com/news/2018-09-crunched-high-intensity-cell-benefits-minutes.html>

| |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|