

Leptospirosis strains identified in Uruguay cattle

September 13 2018



One of the veterinarians, co-author of the article, was photographed while obtaining urine and blood samples from cattle in a farm in Uruguay. Such samples allowed for serologic characterization of animals, and eventual isolation/typing of Leptospira spp. strains. Credit: Buschiazzo, et al. (2018)

Leptospirosis infections, caused by *Leptospira* bacteria, occur in people and animals around the world, but different strains of the bacteria may vary in their ability to cause disease and to jump between species. Now,



researchers reporting in *PLOS Neglected Tropical Diseases* have for the first time described the characteristics of the *Leptospira* variants that infect cattle in Uruguay.

Leptospirosis is most often transmitted to humans through contaminated water, and outbreaks are seen in tropical and subtropical areas after large amounts of rain or floods. Water can initially be contaminated through the <u>urine</u> of infected animals, including rats, <u>cattle</u>, horses, sheep, pigs and dogs. In Uruguay, beef and dairy exports are leading sources of national income.

To determine the identity of the *Leptospira* variants that infect cattle in Uruguay, and whether they are a potential risk for humans, a multicentric consortium was created involving the Institut Pasteur of Montevideo, the Faculty of Medicine (Udelar), the Agricultural Research National Institute (INIA) and the Ministry of Livestock (MGAP). This multidisciplinary team sampled urine and blood from 963 cattle at 48 beef and dairy farms around Uruguay. Additionally, they collected the urine and kidneys from 577 animals from 22 slaughterhouses. Each sample was tested for the presence of *Leptospira* and, if present, for the exact strain.

The researchers found that 20% of all cattle sampled were shedding pathogenic *Leptospira* in their urine, representing a large public health risk. 40 different strains of the bacteria were isolated, uncovering an unexpectedly large variation. The bacteria identified included three rare isolates undetected by normal tests, and two serotypes of the bacteria that matched exactly with those previously isolated from leptospirosis patients.

"This report of local *Leptospira* strains shall improve diagnostic tools and the understanding of leptospirosis epidemiology in South America," the researchers say. "These strains could also be used as new components



within bacterin vaccines to protect against the pathogenic *Leptospira* strains that are actually circulating, a direct measure to reduce the risk of human leptospirosis."

More information: Zarantonelli L, Suanes A, Meny P, Buroni F, Nieves C, et al. (2018) Isolation of pathogenic Leptospira strains from naturally infected cattle in Uruguay reveals high serovar diversity, and uncovers a relevant risk for human leptospirosis. *PLOS Neglected Tropical Diseases* 12(9): e0006694. doi.org/10.1371/journal.pntd.0006694

Provided by Public Library of Science

Citation: Leptospirosis strains identified in Uruguay cattle (2018, September 13) retrieved 23 May 2024 from <u>https://medicalxpress.com/news/2018-09-leptospirosis-strains-uruguay-</u> <u>cattle.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.