

Research examines mechanisms behind cognitive decline in type 2 diabetes

September 19 2018



Credit: CC0 Public Domain

Type 2 diabetes has been linked with an increased risk of cognitive dysfunction and dementia, but the underlying mechanisms are uncertain. In a new *Diabetic Medicine* study, imaging tests revealed that changes in

white matter regions of the brain that are indicative of small vessel disease are associated with decreased processing speed (the the time it takes a person to do a mental task) in people with Type 2 diabetes. Also, higher blood pressure and worse kidney function were related to worse cognition.

The findings suggest that blood vessels in the brain may be involved in the mechanisms that lead to cognitive decline in individuals with Type 2 diabetes.

"Cognitive dysfunction is an emerging target of complications of [diabetes mellitus](#), and we believe that our results provide some clue into the mechanisms behind its development," said lead author Dr. Boris Mankovsky, of the National Medical Academy for Postgraduate Education, in Kiev, Ukraine.

More information: *Diabetic Medicine*,
onlinelibrary.wiley.com/doi/10.1111/dme.13800

Provided by Wiley

Citation: Research examines mechanisms behind cognitive decline in type 2 diabetes (2018, September 19) retrieved 23 April 2024 from
<https://medicalxpress.com/news/2018-09-mechanisms-cognitive-decline-diabetes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--