


Optimizing dopaminergic treatment improves non-motor symptoms

September 24 2018



(modified; 0-276, higher = worse); D, domain. Frequencies of the modified

NMSS in the "off" and the "on" state. Credit: Swedish Research Council

Non-motor symptoms are common in late stage Parkinson's disease (PD) as the frequency and severity of most of these symptoms increase with advancing disease. Optimizing dopaminergic treatment in the most severe stages can affect non-motor symptoms and improve quality of life, report scientists in the *Journal of Parkinson's Disease*.

PD is generally considered a disease that affects movement, but it also involves a large number of non-motor symptoms, which previous research has shown to have a greater impact on health-related quality of life than motor symptoms. Common non-motor symptoms include cognitive impairment, mood-related symptoms such as depression, apathy, sleep/daytime sleepiness, fatigue, and autonomic dysfunction such as urinary urgency, incontinence, and erectile dysfunction. The frequency and severity of most of these symptoms increase with advancing disease. Previous research has shown that the frequency and severity of non-motor symptoms as a whole are the most important predictors of health-related quality of life in [patients](#) with PD.

"Patients in late stage PD—the last four or five years of the disease—are a forgotten group, whose situation we do not know much about," explained lead investigator Per Odin, MD, Ph.D., Professor, Department of Neurology, Lund University, Lund, Sweden, and Chairman, Department of Neurology, Central Hospital, Bremerhaven, Germany. "There is reason to believe that relatively many patients in the late stage of PD may be insufficiently treated. Since the effect of dopaminergic therapy may not be as obvious in the late stage as in earlier disease stages, there is a clear risk for undertreatment. The general aim of the present research is therefore to learn more about this group of patients, to get a basis for improving their situation."

The goal of the present study was to find an optimal pharmacological treatment for patients with late stage PD. Investigators assessed the effect of dopaminergic therapy on non-motor symptoms in 30 patients. Participants were recruited from the southern region of Sweden through neurology departments and the municipality-based health care system. The patients were in "Hoehn and Yahr" stages IV and V; the Hoehn and Yahr scale is a commonly used system for classifying how the symptoms of PD progress.

The dopaminergic effect on non-motor symptomatology was assessed using a Non-Motor Symptoms Scale (NMSS) in the "off" and the "on" state during a standardized L-dopa test and assessing non-motor symptoms in parallel with motor function. Motor symptoms fluctuate between an "on" state, during which the patient experiences a positive response to medication, and an "off" state, during which the patient experiences a re-emergence of the Parkinson symptoms suppressed during the "on" state.

The study found that non-motor symptoms were common and many of the symptoms occurred in more than 80% of the individuals. The highest scores (frequency x severity) were seen within the NMSS domains 3: mood/apathy and 7: urinary in both the "off" and the "on" states. The differences in the NMSS score between the "off" and the "on" state were larger in general for motor responders than for motor non-responders.

The investigators concluded that there is often an L-dopa effect on both motor and non-motor symptoms even in the most severely ill PD patients, and that even if there is not a significant motor response, there was often a non-motor effect, particularly on mood or depressive symptoms. They emphasize the importance of optimizing L-dopa treatment in the late stage of the disease to give patients the best possible quality of life.

"We encourage colleagues who treat PD to pay attention so that the treatment is optimized throughout the disease progression and into the most severe disease stages," commented Dr. Odin. "The knowledge that sufficient dopaminergic treatment may have important effects on both motor and non-motor symptoms in severe PD patients can hopefully help treating physicians improve quality of life for their patients with late [stage](#) PD.

More information: Kristina Rosqvist et al, Dopaminergic Effect on Non-Motor Symptoms in Late Stage Parkinson's Disease, *Journal of Parkinson's Disease* (2018). [DOI: 10.3233/JPD-181380](https://doi.org/10.3233/JPD-181380)

Provided by IOS Press

Citation: Optimizing dopaminergic treatment improves non-motor symptoms (2018, September 24) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-09-optimizing-dopaminergic-treatment-non-motor-symptoms.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--