

Prostate cancer care for older men estimated to cost Medicare \$1.2 billion

September 13 2018



Micrograph showing prostatic acinar adenocarcinoma (the most common form of prostate cancer) Credit: Wikipedia, [CC BY-SA 3.0](https://creativecommons.org/licenses/by-sa/3.0/)

Researchers from the University of North Carolina Lineberger Comprehensive Cancer Center estimate that screening for and treating prostate cancer in men aged 70 years or older, which is not recommended by national guidelines, cost Medicare more than \$1.2 billion over a three year period for each group of men diagnosed in the United States each year.

Published in the journal *JAMA Oncology*, the study examined the costs associated with screening for [prostate cancer](#), including [treatment](#), for three years after diagnosis. They estimated that for men diagnosed in each of 2004, 2005, 2006, and 2007, the total cost for treating and screening for each group would be \$1.2 billion for three years after diagnosis. The study also showed that of the total, \$451 million was spent on men who were diagnosed with prostate cancer that is graded with a Gleason score of 6, which is considered to be low-grade disease and best managed using active surveillance rather than aggressive treatment.

"The tough discussions that happen in health economics are often cases where care is beneficial, but costly. That's a hard trade-off, but this one is actually easier than that," said UNC Lineberger's Justin Trogon, Ph.D., associate professor of health policy management in the UNC Gillings School of Global Public Health, and the study's first author. "This is a scenario where care is probably not beneficial and also costly, and we are putting a dollar figure on just how costly this is."

The researchers said there is a wide recognition that screening for prostate cancer should stop in men at age 70. The U.S. Preventive Services Task Force recommends against [prostate cancer screening](#) in men above age 70, citing, among other reasons, that the harms from screening are "at least moderate" or greater for men in that age group because of risk of false-positives, harms from biopsy, and harms from treatment, which can include [sexual dysfunction](#) and [urinary incontinence](#).

"For [patients](#) who are diagnosed with prostate cancer at an older age, because the cancer is often slow-growing, it is unlikely to be the cause of their mortality," said UNC Lineberger's Ronald C. Chen, MD, MPH, associate professor in the UNC School of Medicine Department of Radiation Oncology and the study's corresponding author. "So, we are

not only spending U.S. health care dollars diagnosing an issue that probably isn't going to be a problem for patients, but when we treat these patients who don't need treatment, they suffer unnecessary side effects like urinary incontinence or sexual dysfunction."

For the retrospective study, researchers used the National Cancer Institute Surveillance, Epidemiology, and End Results (SEER)-Medicare linked claims database to analyze costs related to diagnosis and work-up, treatment, follow-up, and side-effect management for nearly 50,000 men 70 years of age or older who were diagnosed with prostate cancer between 2004 and 2007. They estimated the cost of diagnosing and treating men for three years after diagnosis because that's a typical time frame for patients to receive treatment in.

They found that for men diagnosed in each year, the total three-year cost was \$1.2 billion. For men diagnosed in 2004, for example, their estimated cost to Medicare would be \$1.2 billion across three years. The median per-patient cost within three years after prostate cancer diagnosis was \$14,453, with treatment costs accounting for 73 percent of that total. The researchers estimated a cost savings of \$320 million per year for Medicare if patients who have a Gleason score of 6 choose active surveillance.

"Studies indicate that we are testing for [prostate](#) cancer in people for whom the potential harms outweigh the potential benefits, and then we are treating—actually over treating—people for a cancer that may have little to no impact on their life expectancy," Chen said. "Not following guidelines could be putting people's health at risk, and it is costing the U.S. health care system a lot of money. It would make more sense to put these resources into other areas that could do more good for patients."

Researchers write in the study that foregoing aggressive treatment of [low-risk prostate cancer](#) in older patients, especially those with other health

issues, could prevent harm due to treatment-related side effects, while also providing a savings to Medicare. They suggest that screening men with at least a 10-year life expectancy, and selectively treating those with more aggressive [cancer](#) may be one approach to eliminate unnecessary spending while maximizing patient outcomes.

Provided by UNC Lineberger Comprehensive Cancer Center

Citation: Prostate cancer care for older men estimated to cost Medicare \$1.2 billion (2018, September 13) retrieved 24 April 2024 from <https://medicalxpress.com/news/2018-09-prostate-cancer-older-men-medicare.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.