

New screening tool can improve the quality of life for epilepsy patients with sleep apnea

September 27 2018



The study was led by Martha A. Mulvey (left), a nurse practitioner at University Hospital's department of neurosciences. Co-author Xue Ming is a professor of neurology at Rutgers New Jersey Medical School. Credit: Nick Romanenko / Rutgers University

Rutgers researchers have developed a tool to help neurologists screen for

obstructive sleep apnea in people with epilepsy whose seizures can be magnified by sleep disorders.

The study appears in the journal *Neurology Clinical Practice*.

Although detection and treatment of [obstructive sleep apnea](#) (OSA) can improve [seizure](#) control in some patients with epilepsy, providers have not regularly assessed patients for those [risk factors](#). The researchers developed an electronic health record alert for neurologists to evaluate a patient's need for a sleep study.

This study can determine the necessity for treatment, which can result in improved seizure control, reduction in antiepileptic medications and reduce the risk of sudden unexpected death in epilepsy.

OSA occurs when breathing is interrupted during sleep. The Epilepsy Foundation estimates that approximately 40 percent of people living with epilepsy have a higher prevalence of OSA that contributes to poor [seizure control](#).

"Sleep disorders are common among people living with epilepsy and are under-diagnosed," said lead author Martha A. Mulvey, a nurse practitioner at University Hospital's department of neurosciences. "Sleep and epilepsy have a complex reciprocal relationship. Seizures can often be triggered by low oxygen levels that occur during OSA. Sleep deprivation and the interruption of sleep can therefore increase seizure frequency."

The researchers developed an assessment for identifying OSA consisting of 12 recognized risk factors, which are embedded in the electronic health record. If a patient has at least two risk factors, they are referred for a sleep study. The risk factors include: body mass index greater than 30 kg/m²; snoring; choking or gasping in sleep; unexplained nighttime

awakenings; morning headaches; dry mouth, sore throat or chest tightness upon awakening; undue nighttime urination; decreased memory and concentration; neck circumference greater than 17 inches; excessive daytime sleepiness; undersized or backward displacement of the jaw; and an assessment of the distance from the tongue base to the roof of the mouth.

"It was found that placing this mandatory alert for providers to screen for OSA in the EHR markedly increased the detection of at-risk epilepsy patients who should be referred for a sleep study," said co-author Xue Ming, professor of neurology at Rutgers New Jersey Medical School. "Such screening can lead to early detection and treatment, which will improve the quality of life of patients with epilepsy and OSA."

In cases that were reviewed prior to the alert being placed in the electronic health record, only 7 percent with [epilepsy](#) were referred for sleep studies. Of those who were referred, 56 percent were diagnosed with sleep apnea. Of the 405 patients who were screened for OSA after the alert was placed in the electronic health record, 33 percent had at least two risk factors and were referred for a sleep study. Of the 82 [patients](#) who completed a sleep study, 87 percent showed at least mild sleep apnea.

More information: [DOI: 10.1212/CPJ.0000000000000502](https://doi.org/10.1212/CPJ.0000000000000502)

Provided by Rutgers University

Citation: New screening tool can improve the quality of life for epilepsy patients with sleep apnea (2018, September 27) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-09-screening-tool-quality-life-epilepsy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.