

Sleepiness common among black women, may be linked to high blood pressure

September 6 2018

Poor sleep habits may be related to low levels of physical activity, high blood pressure and obesity among black women, according to new research presented at the American Heart Association's Joint Hypertension 2018 Scientific Sessions, an annual conference focused on recent advances in hypertension research.

In a study of overweight and obese black women, researchers set out to determine whether there's a connection between sleep, [blood pressure](#) and levels of [physical activity](#). Sleepiness was determined using a questionnaire that asks whether a person is likely to "doze" during certain daily activities, and their likelihood of napping if given the opportunity. The women were 50 years old on average.

About half of the 361 women studied reported sleeping less than 7 hours a night. In addition, 62 percent reported daytime sleepiness (35 percent at "high normal" and 27 percent at "excessive"). The study found these women to have a greater incidence of limited sleep and excessive daytime sleepiness compared to the general population.

Women who reported high levels of daytime sleepiness also reported much less vigorous and moderate physical activity and had higher blood pressure compared to those who said they had normal levels of [daytime sleepiness](#).

These findings suggest that efforts to combat obesity and [high blood pressure](#) among this population should address sleep quality, the authors

said.

Provided by American Heart Association

Citation: Sleepiness common among black women, may be linked to high blood pressure (2018, September 6) retrieved 3 May 2024 from <https://medicalxpress.com/news/2018-09-sleepiness-common-black-women-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.