


# Researchers uncover previously unknown dementia, Alzheimer's risk factor

September 21 2018, by Orli Belman



Structure of the human mitochondrial genome. Credit: Wikipedia/CC BY-SA 3.0

New research from USC has uncovered a previously unknown genetic risk factor for Alzheimer's disease and related dementias. The study provides insights on how these conditions, and other diseases of aging,

might one day be treated and prevented.

The research from the Cohen Lab at the USC Leonard Davis School of Gerontology sheds new light on the protective role of a naturally occurring [mitochondrial](#) peptide, known as humanin. Amounts of the peptide decrease with age, leading researchers to believe that humanin levels play an important function in the aging process and the onset of diseases linked to older age.

"Because of the beneficial effects of humanin, a decrease in circulating levels could lead to an increase in several different diseases of aging, particularly in [dementia](#)," said senior author Pinchas Cohen, dean of the USC Leonard Davis School and one of three researchers to independently discover the existence of humanin 15 years ago.

The study, led by Kelvin Yen of the USC Leonard Davis School, appears online on Sept. 21 in the *Nature*-published journal *Scientific Reports*. Among the findings, the researchers discovered a significant difference between the circulating levels of humanin in African-Americans, who are more impacted by Alzheimer's disease and other diseases of aging, as compared to Caucasians.

"We have now discovered a novel underlying biological factor that may be contributing to this health disparity," said Yen, a research assistant professor of gerontology.

Because humanin is encoded within the [mitochondrial genome](#), the research team examined the mitochondrial DNA for common genetic variations known as SNPs (pronounced snips) that could explain the differences between humanin levels. According to the paper, after sequencing the entire mitochondrial genome of all samples to determine if any SNPs correlated with humanin levels, they identified a genetic variation that was associated with a 14 percent decrease in circulating

humanin levels.

This provides the first evidence that a variation in the sequence of a mitochondrial peptide is associated with a change in the level of [peptides](#) and the first conclusive demonstration that mitochondrial peptides are encoded in and regulated through mitochondrial DNA, Cohen said.

The team subsequently examined the effect of this SNP in a separate large cohort of participants from the Health and Retirement Study, a longitudinal study of approximately 20,000 individuals over the age of 50 in the United States, of which more than 12,500 individuals consented to DNA analysis. Using this data set, the researchers report finding that the unfavorable version of this SNP, associated with low levels of the humanin peptide, is also associated with accelerated cognitive aging, providing the first demonstration linking a humanin SNP in the mitochondria to cognitive decline in people.

The paper also shows that in mice, injections of humanin delay cognitive decline associated with aging, proposing a possible therapeutic role for humanin-related drugs. The mechanism through which humanin acts to mediate these effects involves suppression of inflammation systemically, as well as specifically in the brain.

Provided by University of Southern California

Citation: Researchers uncover previously unknown dementia, Alzheimer's risk factor (2018, September 21) retrieved 23 May 2024 from <https://medicalxpress.com/news/2018-09-uncover-previously-unknown-dementia-alzheimer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--