

How weight loss is linked to future health for older adults

September 4 2018

Studies describing the effects of weight loss on health rarely consider age. However, weight loss during middle age likely has different effects on your health than does weight loss when you're 65-years-old or older—especially when you're older than 85.

Although some studies have found that [weight loss](#) in older adults is generally linked to an increase in illness and death, researchers say that these studies were either too short or were based on information that may have been interpreted incorrectly.

However, one study about fractures and osteoporosis (a medical condition in which bones become thin, lose density, and become increasingly fragile) looked specifically at health and [weight](#) for [women](#) who were over age 65. Reviewing more than 20 years' worth of data for study participants, the team of researchers responsible for this study had the chance to examine links between long-term weight gain/[loss](#) and health. Their findings were published in the Journal of the American Geriatrics Society.

In their new study, the research team evaluated information from the Study of Osteoporotic Fractures. The research team theorized that women with greater weight loss, greater variability in their weight, and/or abrupt declines in weight would be less able to function physically at year 20, and would be more likely to experience poorer health outcomes one to five years after year 20. This theory was based in part on an earlier, related study by the same research team also making use of

the Study of Osteoporotic Fractures. In that earlier work, the researchers discovered that the rate of weight loss over 20 years was linked to developing mild cognitive impairment or dementia in women surviving past age 80. (This was not the case for participants with sudden weight loss or changes in weight).

The current study revealed that every 22 pounds of weight loss over 20 years was linked to a 23 percent increased [risk](#) of death and a 52 percent increased risk of hip fracture.

The team also said that women with moderate weight loss (20 or more pounds) over 20 years had a 74 percent increased risk of death. Their risk for hip fracture increased nearly three times, compared to women who had not lost weight. They were nearly four times more likely to have poor physical function after 20 years, compared to women with no weight loss.

Even women who had lost a small amount of weight (less than 20 pounds) over 20 years had an increased risk of death, but no increased risk of [hip fracture](#) or of poor physical function.

However, the researchers found no link between weight loss and chances for experiencing two or more falls during approximately 18 months of follow-up.

Weight variability and abrupt weight loss were not associated with poor health outcomes, such as falls, fractures, and death. However, those with the most weight variability over 20 years were two times more likely to have poor scores for measures of physical function.

As women age, they risk weight loss because of changes in senses of taste and smell, poorer digestion, and difficulty absorbing nutrients. In addition, other challenges such as loneliness, being in a long-term care

facility, having mental health problems such as depression, and/or having limited ability to get around independently can lead to weight loss, said the researchers. "Our findings suggest that weight loss may contribute to the process of health decline," said Dr. Erin LeBlanc, lead author of the study at Kaiser Permanente's Center for Health Research, Portland, Oregon.

The researchers added: "Our results suggest long-term weight loss in older women may be a marker for increased risk of poor health outcomes. Therefore, we should pay attention to women who have survived into their 80s and 90s who have experienced moderate weight loss, regardless of whether there was an abrupt weight decline." Looking closely at women's nutrition, as well as social, environmental, and physical factors impacting well-being also could help preserve [health](#) and physical function into old age. However, additional research is needed, the researchers concluded.

More information: Erin S. LeBlanc et al, Long-Term Weight Trajectory and Risk of Hip Fracture, Falls, Impaired Physical Function, and Death, *Journal of the American Geriatrics Society* (2018). [DOI: 10.1111/jgs.15532](#)

Provided by American Geriatrics Society

Citation: How weight loss is linked to future health for older adults (2018, September 4) retrieved 20 April 2024 from <https://medicalxpress.com/news/2018-09-weight-loss-linked-future-health.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--