


Abdominal obesity common among women in rural Indonesia

October 25 2018



Cahya Utamie Pujilestari, PhD student at the Department of Public Health and Clinical Medicine at Umeå University. Credit: Lena Mustonen

Abdominal obesity is seven times more common among women than among men in Indonesia's rural area. This is shown in a new doctoral thesis at Umeå University, Sweden.

"If preventive measures are not taken, the abdominal obesity will increase among the poorest with severe health effects as a consequence," says Cahya Utamie Pujilestari, Ph.D. student at the Department of Public Health and Clinical Medicine at Umeå University.

In her [doctoral thesis](#), Cahya Utamie Pujilestari has analysed abdominal obesity from a socio-economic and gender equality perspective. The thesis is based on four studies with a total of 15,500 people in the Purworejo district, which consists predominantly of rural areas in the Central Java Province.

Abdominal [obesity](#) was in itself more common in the more prosperous part of the population. On the other hand, disability and death as a result of [abdominal obesity](#) was more common among the poorer population.

"Preventive efforts should be directed at socially vulnerable women which is the group at greatest risk of suffering," says Cahya Utamie Pujilestari.

More information: Abdominal obesity among older population in Indonesia: socioeconomic and gender inequality, pattern and impacts on disability and death. umu.diva-portal.org/smash/record.jsf?pid=diva2%3A1252590&dswid=newPopUp

Provided by Umea University

Citation: Abdominal obesity common among women in rural Indonesia (2018, October 25) retrieved 24 May 2024 from <https://medicalxpress.com/news/2018-10-abdominal-obesity-common-women-rural.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.