


# Half the brain encodes both arm movements

October 8 2018




Patients implanted with electrocorticography arrays completed a 3D center-out reaching task. Electrode locations were based upon the clinical requirements of each patient and were localized to an atlas brain for display (A). B. Patients were seated in the semi-recumbent position and completed reaching movements from the center to the corners of a 50cm physical cube based upon cues from LED lights located at each target while hand positions and ECoG signals were simultaneously recorded. Each patient was implanted with electrodes in a single cortical hemisphere and performed the task with the arm contralateral (C) and ipsilateral (D) to the electrode array in separate recording sessions. Credit: Bundy et al., *JNeurosci* (2018)

Individual arm movements are represented by neural activity in both the left and right hemispheres of the brain, according to a study of epilepsy patients published in *JNeurosci*. This finding suggests the unaffected hemisphere in stroke could be harnessed to restore limb function on the same side of the body by controlling a brain-computer interface.

The right side of the brain is understood to control the left side of the body, and vice versa. Recent evidence, however, supports a connection between the same side of the brain and body during [limb movement](#).

Eric Leuthardt, David Bundy, and colleagues explored brain activity during such ipsilateral movements during a reaching task in four [epilepsy patients](#) whose condition enabled invasive monitoring of their brains through implanted electrodes. Using a machine learning algorithm, the researchers demonstrate successful decoding of speed, velocity, and position information of both left and right arm movements regardless of the location of the electrodes.

In addition to advancing our understanding of how the brain controls the body, these results could inform the development of more effective rehabilitation strategies following brain injury.



In the study a patient implanted with electrodes only on the left side of the brain was asked to make movements to 8 targets in 3D space with both their right and left arms. Using recordings from these electrodes, the authors were able to predict the hand speed, direction, and position for both arms showing that movements of both arms are encoded on one side of the brain. Credit: David Bundy and Eric Leuthardt

**More information:** Unilateral, Three-dimensional Arm Movement Kinematics are Encoded in Ipsilateral Human Cortex, *JNeurosci* (2018). DOI: [10.1523/JNEUROSCI.0015-18.2018](https://doi.org/10.1523/JNEUROSCI.0015-18.2018)

Provided by Society for Neuroscience

Citation: Half the brain encodes both arm movements (2018, October 8) retrieved 10 April 2024 from <https://medicalxpress.com/news/2018-10-brain-encodes-arm-movements.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.