

Study shows childhood obesity the major risk factor for serious hip disease

October 22 2018

New research suggests that rising childhood obesity rates are causing more adolescents to develop a debilitating hip disease requiring urgent surgery.

Slipped Capital Femoral Epiphysis (SCFE) is a disease that occurs amongst teenagers, affecting 1 in 2,000 [children](#). The hip deforms, and sometimes completely collapses, causing pain and life-long disability. Early recognition and treatment of the disease is important to minimise the severity. Nevertheless, SCFE is one of the most common reasons for a hip replacement in young adults, and sometimes even children. Surgeons have long believed [obesity](#) may be responsible, though there was no substantial proof until now.

The study, "Childhood Obesity and Slipped Capital Femoral Epiphysis" (online Oct. 22), published in *Pediatrics*, was conducted by researchers from the University of Liverpool, University of Oxford, University of Aberdeen and Alder Hey Children's Hospital.

The study looked at body mass index (BMI) for nearly 600,000 children in Scotland, where BMI measurements are collected through routine school health screening. Using this very large cohort of children, the researchers showed that children obese at 5 years old, had a 75% likelihood of remaining obese when they were 12 years old. Most crucially, children with severe obesity at 5 years old, had almost 20 times the subsequent risk of developing SCFE than a thin child. The greater the BMI of the child, the greater the risk of SCFE.

Daniel Perry, Children's Orthopaedic Surgeon at Alder Hey Hospital and NIHR Clinician Scientist said: "Surgeons have long suspected that [childhood obesity](#) was the cause of this disease, and these results make it very clear. It is important that doctors who treat children are aware of SCFE, especially amongst children with obesity. Identifying SCFE early means children typically only need relatively simple surgery, however children identified later often require high-risk reconstructive surgery".

"High rates of Childhood Obesity are of major national and international concern. SCFE has a profound affect on the quality of lives of adolescents. The link with obesity is striking—there are few other diseases in children that occur directly as a consequence of obesity, and this disease causes lifelong problems for the child", adds Professor Steve Turner, Professor of Paediatrics at the University of Aberdeen.

More information: *Pediatrics* (2018). [DOI: 10.1542/peds.2018-1067](https://doi.org/10.1542/peds.2018-1067)

Provided by University of Liverpool

Citation: Study shows childhood obesity the major risk factor for serious hip disease (2018, October 22) retrieved 27 April 2024 from <https://medicalxpress.com/news/2018-10-childhood-obesity-major-factor-hip.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--