

Study shows diet and weight may affect response to bipolar disorder treatment

October 7 2018

Data from a clinical trial has shown that how people respond to treatment for Bipolar Disorder may be influenced by their weight and the overall quality of their diet, including whether they are eating a diet high in foods thought to contribute to general inflammation. These are early results, but if replicated may mean that treatment of some mental health problems could benefit from the inclusion of dietary advice. This is presented at the ECNP Conference in Barcelona.

Bipolar Disorder (which used to be called 'manic depression') is characterised by episodes of mood swings, between being very up or very down with periods in between the two extremes. The fact that there are two opposite sets of symptoms means that finding an effective [treatment](#) is difficult. While current medications are useful, they are better at targeting mania symptoms (the 'up' phase), leaving a lack of [effective treatment](#) for people experiencing depressive episodes. Now a group of Australian, German and American scientists have shown those who have a high quality diet, a less inflammatory diet, and/or a low BMI (Body Mass Index) may respond better to an add-on nutraceutical treatment provided as part of a clinical trial.

"If we can confirm these results, then it's good news for people with Bipolar Disorder, as there is a great need for better treatments for the depressive phase of Bipolar Disorder" said lead researcher Melanie Ashton of Deakin University in Australia.

A total of 133 participants were randomly assigned to take a

combination of nutraceuticals (compounds derived from foods such as vitamins or minerals that treat or prevent a disease or disorder) including the anti-inflammatory amino acid n-acetylcysteine (NAC), or NAC alone, or a placebo (a dummy pill) for 16 weeks. Participants received the study medication in addition to any stable treatments they were already receiving. Researchers measured BMI at the beginning of the study, and then measured depression and how a person is able to function in their day to day life. Researchers also rated whether a participant was improving and, if so, how much, over the next 20 weeks. Participants filled in a questionnaire about what they usually eat over the year and researchers calculated a diet quality score, where good diets included a healthy diet with lots of fruit and vegetables, whereas poorer-quality diets had more saturated fat, refined carbohydrates and alcohol. These types of diets were then categorised as either anti-inflammatory or pro-inflammatory based on foods that affect inflammation.

Melanie Ashton continued, "We found that people who had a better-quality diet, a diet with anti-inflammatory properties, or a lower BMI, showed better response to add-on nutraceutical treatment than did those who reported a low-quality diet, or a diet including foods that promote inflammation, or who were overweight.

What this means, if these results can be repeated in a larger trial, is that treatment for Bipolar Disorder would need to take into account what a person eats and their weight.

There are some points we need to note about this study. This is a randomised, controlled trial, but what we found were exploratory outcomes; in other words, it wasn't the main result that we were testing. Our result is statistically significant, but because the study wasn't specifically designed to test the effect of diet quality,

inflammatory diets and BMI on drug response in general, it is necessary

to see the work replicated in a larger study before any firm conclusions can be formed".

Commenting, Professor Eduard Vieta (Barcelona) said:

"This is interesting work, which holds out the possibility that patients with Bipolar Disorder may benefit from a balanced [diet](#). However, it is an early study, and we need more research before we can think whether this might affect clinical practice".

Professor Vieta was not involved in this work, it is an independent comment.

Provided by European College of Neuropsychopharmacology

Citation: Study shows diet and weight may affect response to bipolar disorder treatment (2018, October 7) retrieved 11 May 2024 from <https://medicalxpress.com/news/2018-10-diet-weight-affect-response-bipolar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.