

Discharged mental health patients 'at greater risk of dying'

October 3 2018, by Michael Addelman



Credit: CC0 Public Domain

Mental health patients are at much greater risk of dying from unnatural causes following their first discharge from inpatient care than the rest of the population, according to new research.

In the most comprehensive study of this topic ever carried out, The University of Manchester team say though danger to patients is still relatively low, community services and GPs must be aware of the critical period in the first few weeks of discharge—when they are most at risk.

Though the report drew on data derived the entire population of Denmark, its results are relevant to the UK, which has a similar population and health system, say authors Professor Roger Webb and Ph.D. student Florian Walter.

According to the team, the most common threat to life among patients after discharge is fatal poisoning compared to accidents among the [general population](#).

The research was funded by Medical Research Council, European Research Council in partnership with Centre for Integrated Register-based Research and National Centre for Register-based Research at Aarhus University. It is published in the *Journal of Clinical Psychiatry* today.

The analysis compared over 1.6 million Danes with a subset of 47,000 people who were discharged from their first inpatient stay between 1982 and 1996.

Utilising national administrative datasets, the research team were able to track the group of patients, who averaged age 25 when they were discharged until they died or 31 December 2011.

They looked at method of suicide, types of accidental death, types of fatal poisoning and death by homicide.

Though there is a strong association between hospital discharge and unnatural death, there is no evidence of a causal link between the two,

say the researchers.

According to the research, suicide was 32 times more likely after discharge: 171.7 people in every 100,000 died after leaving hospital, compared to 5.3 in every 100,000 for the general population

Intentional self-poisoning was 41 times more likely: 57.1 people in every 100,000 died after hospital discharge, compared to 1.3 in every 100,000 for the general population

And death from fatal psychotropic drugs poisoning , irrespective of intent was over 90 times more likely: 25.6 people in every 100,000 after [hospital discharge](#), compared to 0.2 in every 100,000 for the general population.

Professor Webb said: "When we look at detailed causes of death, each one is markedly raised whether that's traffic accidents or poisoning.

"So it's clear this group of people is more vulnerable to dying from a variety of different unnatural causes after discharge, although the risks for each of these causes are nonetheless relatively low in absolute terms.

"However, it's important to point out hospital is not a cause of unnatural death as these patients are a young cohort already from high risk groups; therefore the risk of dying could be even higher without hospitalisation."

Florian Walter said: "This is the first study to look at the full range unnatural causes of death among [mental health patients](#) after discharge from hospital ever carried out.

"The Danish system is ideal for this sort of research because Danes are given a unique identification number which provides data that are entered into several national registers."

A previous study by Walter, published in *JAMA Psychiatry* in 2017, found that patients are most at risk in the first month after discharge.

He added: "The overarching message is that transition from [hospital](#) to community is riskier for these people, especially during the first few weeks after discharge.

"Therefore mental health services need to develop broad strategies that include enhanced post-[discharge](#) surveillance. These strategies must account for the fact that discharged patients have higher risks of dying unnaturally in lots of different ways.

"We also think better liaison between inpatient and community services is important so that services can be more vigilant about patients taking their medication correctly at the doses prescribed."

The paper, "Suicide methods and specific types of accidental death and fatal poisoning among discharged psychiatric patients: a national cohort study," is published in the *Journal of Clinical Psychiatry*.

More information: Florian Walter et al. Premature Mortality Among Patients Recently Discharged From Their First Inpatient Psychiatric Treatment, *JAMA Psychiatry* (2017). [DOI: 10.1001/jamapsychiatry.2017.0071](#)

Florian Walter et al. Suicide methods and specific types of accidental death and fatal poisoning among discharged psychiatric patients: a national cohort study, *Journal of Clinical Psychiatry*, 04.04.2018

Provided by University of Manchester

Citation: Discharged mental health patients 'at greater risk of dying' (2018, October 3) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2018-10-discharged-mental-health-patients-greater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.