

# Domestic refrigerators may pose risk to insulin quality

October 3 2018

---

New research being presented at this year's European Association for the Study of Diabetes (EASD) Annual Meeting in Berlin, Germany (1-5 October), suggests that insulin is often stored at the wrong temperature in patients' fridges at home, which could affect its potency.

Many injectable drugs and vaccines are highly sensitive to heat and cold and can perish if their temperature shifts a few degrees. To prevent loss of effectiveness, [insulin](#) must stay between 2-8°C/36-46°F in the refrigerator or 2-30°C/30-86°F when carried about the person in a pen or vial.

Individuals with [diabetes](#) often store insulin at home for several months before they use it, but little is known about how storage in domestic fridges impacts insulin quality.

To investigate how often insulin is stored outside the manufacturer's recommended temperature range, Dr. Katarina Braune from Charité - Universitaetsmedizin Berlin in Germany in collaboration with Professor Lutz Heinemann (Science & Co) and the digital health company MedAngel BV monitored the temperature of insulin formulations stored in fridges at home and carried as a spare.

Between November 2016 and February 2018, 388 diabetes patients living in the USA and the EU placed temperature sensors (MedAngel ONE, <http://www.medangel.co>) either next to their insulin in the fridge and or their diabetes bag.

Temperature data were automatically measured every 3 minutes (up to 480 times a day) before being sent to an app and recorded on a secure database. Temperature data were recorded for an average of 49 days.

Analysis of 400 temperature logs (230 for refrigerated and 170 carried insulin) revealed that 315 (79%) contained deviations from the recommended temperature range.

On average, insulin stored in the fridge was out of the recommended temperature range 11% of the time (equivalent to 2 hours and 34 mins a day). In contrast, insulin carried by patients was only outside recommendations for around 8 minutes a day.

Importantly, freezing was an even bigger issue, with 66 sensors (17%) measuring temperatures below 0°C (equivalent to 3 hours a month on average).

"Many people with diabetes are unwittingly storing their insulin wrong because of fluctuating temperatures in domestic refrigerators", says Dr. Braune.

"When storing your insulin in the fridge at home, always use a thermometer to check the temperature. Long-term storage conditions of insulin are known to have an impact on its blood-glucose lowering effect. For people living with insulin-dependent diabetes who take insulin several times a day via injections or continuously administer insulin with a pump, precise dosing is essential to achieve optimal therapeutic outcomes. Even gradual loss of potency introduces unnecessary variability in dosing. More research is needed to examine the extent to which [temperature](#) deviations during domestic storage affect insulin efficacy and patient outcomes."

Provided by Diabetologia

Citation: Domestic refrigerators may pose risk to insulin quality (2018, October 3) retrieved 7 May 2024 from <https://medicalxpress.com/news/2018-10-domestic-refrigerators-pose-insulin-quality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.