

Experimental vaccine may reduce post-stroke blood clot risk

October 29 2018



The brain has about 100 billion cells called neurons. It's made up of distinct parts, that developed through human evolution. Credit: American Heart Association

A vaccine may one day be able to replace oral blood thinners to reduce the risk of secondary strokes caused by blood clots, without increasing the risk of serious bleeding or triggering an autoimmune response, according to new research in the American Heart Association's journal

Hypertension.

People who have had a stroke caused by a blood clot (ischemic strokes) often need to take medications that make their blood less likely to clot, which helps prevent another stroke.

Japanese researchers successfully tested an experimental [vaccine](#) in mice and found that it provided protection against [blood clots](#) for more than two months without increasing the risk of bleeding or causing an autoimmune response. The lack of an [autoimmune response](#) is important, because it means the mice's immune system did not perceive the vaccine as an "intruder" that needed to be attacked, which would have caused a reaction to the vaccine.

The vaccine, S100A9, inhibits [blood clot formation](#) and, during the study, protected the arteries of treated mice from forming new clots for more than two months, and additionally, worked as well as the oral blood thinner clopidogrel in a major artery, according to Hironori Nakagami, M.D., Ph.D., study co-author and professor at Osaka University, in Japan.

Developing a vaccine to replace and/or compliment daily, oral medications might save many lives and help prevent both secondary strokes and possibly heart attacks, according to Nakagami.




Illustration showing a blood flow block in the brain during an ischemic stroke.
Credit: American Heart Association

"Many stroke patients don't take their blood thinning drugs as prescribed, which makes it more likely they will have another stroke. This vaccine might one day help solve this issue since it would only need to be injected periodically," Nakagami said.

"We are continuing our research in hopes of being able to start clinical trials between five and ten years from now, but there are differences between mice and humans in how the vaccine will be recognized by the immune system," he said. "We should be able to overcome such problems and believe this vaccine provides a very promising strategy in secondary prevention of stroke."

More information: Tomohiro Kawano et al. Therapeutic Vaccine Against S100A9 (S100 Calcium-Binding Protein A9) Inhibits Thrombosis Without Increasing the Risk of Bleeding in Ischemic Stroke in Mice. *Hypertension*. 2018;0:HYPERTEENSIONAHA.118.11316

Provided by American Heart Association

Citation: Experimental vaccine may reduce post-stroke blood clot risk (2018, October 29)
retrieved 4 May 2024 from
https://medicalxpress.com/news/2018-10-experimental-vaccine-post-stroke-blood-clot_1.html

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--