


Father's nicotine use can cause cognitive problems in children and grandchildren

October 16 2018



A photomicrograph of a histological section of a mouse testis processed for immunohistochemistry for methylated DNA. Spermatozoa at different stages of maturation are labeled with the methylated DNA marker (arrow). The data show that nicotine increases DNA methylation in the spermatozoa. Inset shows isolated mature spermatozoa labeled with an immunohistochemical marker for the dopamine D2 receptor (white arrow). Data show that nicotine produces significant changes in DNA methylation in the promoter region of the D2 receptor gene in the spermatozo. Credit: Pradeep G. Bhide

A father's exposure to nicotine may cause cognitive deficits in his children and even grandchildren, according to a study in mice publishing on October 16 in the open-access journal *PLOS Biology* by Pradeep Bhide of Florida State University in Tallahassee and colleagues. The effect, which was not caused by direct secondhand exposure, may be due to epigenetic changes in key genes in the father's sperm.

Exposure of mothers to nicotine and other components of cigarette smoke is recognized as a significant risk factor for behavioral disorders, including [attention deficit hyperactivity](#) disorder, (or ADHD) in multiple generations of descendants. Whether the same applies to fathers has been less clear, in part because in human studies it has been difficult to separate genetic factors (such as a genetic predisposition to ADHD) from environmental factors, such as direct exposure to cigarette smoke.

To overcome this difficulty, Deirdre McCarthy, Pradeep Bhide and colleagues exposed male mice to low-dose nicotine in their drinking water during the stage of life in which the mice produce sperm. They then bred these mice with females that had never been exposed to nicotine. While the fathers were behaviorally normal, both sexes of offspring displayed hyperactivity, attention [deficit](#), and cognitive inflexibility. When female (but not male) mice from this generation were bred with nicotine-naïve mates, male offspring displayed fewer, but still significant, deficits in cognitive flexibility. Analysis of spermatozoa from the original nicotine-exposed males indicated that promoter regions of multiple genes had been epigenetically modified, including the dopamine D2 gene, critical for brain development and learning, suggesting that these modifications likely contributed to the cognitive deficits in the descendants.

Nicotine and [cigarette smoke](#) have been previously shown to cause widespread [epigenetic changes](#), Bhide said. "The fact that men smoke more than women makes the effects in males especially important from

a public health perspective. Our findings underscore the need for more research on the effects of smoking by the father, rather than just the mother, on the health of their children."

More information: McCarthy DM, Morgan TJ Jr, Lowe SE, Williamson MJ, Spencer TJ, Biederman J, et al. (2018) Nicotine exposure of male mice produces behavioral impairment in multiple generations of descendants. *PLoS Biol* 16(10): e2006497. doi.org/10.1371/journal.pbio.2006497

Provided by Public Library of Science

Citation: Father's nicotine use can cause cognitive problems in children and grandchildren (2018, October 16) retrieved 20 March 2024 from <https://medicalxpress.com/news/2018-10-father-nicotine-cognitive-problems-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
