

Poor oral health linked to higher blood pressure, worse blood pressure control

October 22 2018



Close up of dental exam with mirror. Credit: copyright American Heart Association

People with high blood pressure taking medication for their condition are more likely to benefit from the therapy if they have good oral health,

according to new research in the American Heart Association's journal *Hypertension*.

Findings of the analysis, based on a review of medical and dental exam records of more than 3,600 people with [high blood pressure](#), reveal that those with healthier gums have lower [blood pressure](#) and responded better to blood pressure-lowering medications, compared with individuals who have gum [disease](#), a condition known as periodontitis. Specifically, people with periodontal disease were 20 percent less likely to reach healthy blood pressure ranges, compared with patients in good oral health.

Considering the findings, the researchers say patients with periodontal disease may warrant closer blood pressure monitoring, while those diagnosed with hypertension, or persistently elevated blood pressure, might benefit from a referral to a dentist.

"Physicians should pay close attention to patients' oral health, particularly those receiving treatment for hypertension, and urge those with signs of periodontal disease to seek dental care," Pietropaoli said. "Likewise, dental health professionals should be aware that oral health is indispensable to overall physiological health, including cardiovascular status," said study lead investigator Davide Pietropaoli, D.D.S., Ph.D., of the University of L'Aquila in Italy.

The target blood pressure range for people with hypertension is less than 130/80 mmHg according to the latest recommendations from the American Heart Association/American College of Cardiology. In the study, patients with severe periodontitis had systolic pressure that was, on average, 3 mmHg higher than those with good oral health. Systolic pressure, the upper number in a blood pressure reading, indicates the pressure of blood against the walls of the arteries. While seemingly small, the 3mmHg difference is similar to the reduction in blood

pressure that can be achieved by reducing salt intake by 6 grams per day (equal to a teaspoon of salt, or 2.4 grams of sodium), the researchers said.

The presence of periodontal disease widened the gap even farther, up to 7 mmHg, among people with untreated hypertension, the study found. Blood-pressure medication narrowed the gap, down to 3 mmHg, but did not completely eliminate it, suggesting that periodontal disease may interfere with the effectiveness of blood pressure therapy.

"Patients with high blood pressure and the clinicians who care for them should be aware that good oral [health](#) may be just as important in controlling the condition as are several lifestyle interventions known to help control blood pressure, such as a low-salt diet, regular exercise and weight control," Pietropaoli said.

While the study was not designed to clarify exactly how [periodontal disease](#) interferes with [blood pressure](#) treatment, the researchers say their results are consistent with previous research that links low-grade oral inflammation with blood vessel damage and cardiovascular risk.

More information: *Hypertension* (2018). [DOI: 10.1161/HYPERTENSIONAHA.118.11528](#)

Provided by American Heart Association

Citation: Poor oral health linked to higher blood pressure, worse blood pressure control (2018, October 22) retrieved 4 May 2024 from <https://medicalxpress.com/news/2018-10-poor-oral-health-linked-higher.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.