

PROs indicate brachytherapy alone is the superior treatment

October 22 2018

Patient-Reported Outcomes (PROs) from the NRG Oncology trial RTOG 0232 comparing a combined treatment of external beam therapy and brachytherapy (EBT+B) to transperineal interstitial permanent brachytherapy (B) alone indicate a significantly different clinician and patient-reported late toxicity profile between arms despite similarities in progression-free survival results. This abstract was presented at the American Society for Radiation Oncology (ASTRO) annual meeting and was awarded as a Best of ASTRO presentation for 2018.

NRG-RTOG 0232 evaluated the differences between progression-free survival and patient-reported outcomes on two separate [treatment](#) arms. The first arm was treated with combined partial pelvis EBT+B, whereas the second was treated with B alone. Men with prostate carcinoma were randomly assigned to receive treatment on one of the two arms. Researchers measured changes from baseline to 4 and 24 months using expanded prostate index composite (EPIC) and assessed three prostate cancer-specific PRO domains that included bowel, urinary, and sexual.

"Although progression-free survival was similar on both treatment arms, the addition of external beam therapy yielded a higher percentage of clinician-reported late grade 3 or greater toxicities. The patient-reported data confirms that brachytherapy alone is the superior treatment for this for men with intermediate risk prostate cancer, with less patient-reported side effects.

Brachytherapy alone would also be the most cost-efficient treatment

option for patients," stated Deborah W. Bruner, RN, Ph.D., FANN, the abstract's lead author and the Emory University Senior Vice President for Research.

At 24 months following their treatment, researchers found that the men on the EBT + B arm reported poorer urinary, bowel, and sexual PROs and much greater toxicities than the B alone arm. Late grade 3 or greater clinical reported toxicities were 13% for the EBT+B arm and 7% for the B alone arm. Late grade 3 or greater bladder/genitourinary toxicity, the major symptom of concern, was underreported by clinician-report compared to patient-report. Clinicians reported late grade 3 or greater bladder/genitourinary toxicities of 8% for the EBT+B arm and 3% for the B alone arm compared to patient reports of 34% bladder/genitourinary toxicities for the EBT+B arm and 23% for the B alone arm, further demonstrating the critical role patient-reported outcomes play in assessing treatment efficacy and clinically relevant symptoms.

Provided by NRG Oncology

Citation: PROs indicate brachytherapy alone is the superior treatment (2018, October 22) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2018-10-pros-brachytherapy-superior-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--