

## Weight loss success linked with active selfcontrol regions of the brain

October 18 2018



Credit: CC0 Public Domain

New research suggests that higher-level brain functions have a major role in losing weight. In a study among 24 participants at a weight-loss clinic, those who achieved greatest success in terms of weight loss



demonstrated more activity in the brain regions of the lateral prefrontal cortex associated with self-control. The results of the study were published in *Cell Metabolism* on October 18.

"What we found is that in humans, the control of body weight is dependent largely on the areas of the <u>brain</u> involved in <u>self-control</u> and self-regulation," says Alain Dagher (@alain\_dagher), of the Montreal Neurological Institute and Hospital in Canada. "That area of the brain has the ability to take into account long-term information, such as the desire to be healthy, in order to control immediate desires."

Two hormones, leptin and ghrelin, are known to trigger the body to eat in a <u>weight-loss</u> setting. Previous research confirms that these hormone levels change rapidly when weight is shed. "Everybody who loses weight sees this change in leptin and ghrelin," says Dagher. "It is just that some people, for reasons we do not know, are able to maintain their selfregulation in the face of that signal."

To assess the roles these hormones and self-control have in achieving weight loss, the researchers studied 24 subjects from a weight-loss clinic. Prior to starting a standard 1,200 kcal/day weight-loss diet, all participants received a functional MRI study (fMRI) of the brain, which assessed regions including the lateral prefrontal cortex, which is linked with self-regulation, and the ventral <u>medial prefrontal cortex</u>, a brain area involved in motivation, desire, and value.

Subjects were shown pictures of appetizing foods as well as control pictures of scenery. The researchers compared the brain activity response to the food pictures, particularly the high-calorie food pictures, for each subject at baseline, one month, and three months. "When we show pictures of appetizing foods, the ventral medial prefrontal cortex area becomes more active on fMRI," Dagher says.



During the study, researchers noted that at one month and three months, the signal from the ventral prefrontal cortex went down, and it declined the most in people who were more successful at losing weight. Additionally, the lateral prefrontal <u>cortex</u> signal involved in self-control increased throughout the study.

"In the fMRI, the self-control area increased its activity and the value area decreased its activity," says Dagher. "And the amount of change was predictive of successful weight loss." While all participants lost weight, those who achieved the greatest weight loss had fMRI levels indicating a better ability to self-control. And, at the end of the 3-month study, the hormones ghrelin and leptin were starting to return to baseline, suggesting that a new set point was achieved.

"These results suggest that weight loss treatments that increase selfcontrol, such as cognitive behavioral therapy, may be helpful, particularly when stress is involved in leading to overeating," he says. "Stress disrupts the <u>lateral prefrontal cortex</u> control mechanism, but you may be able train people to seek a different strategy."

**More information:** Neseliler, et al. "Neurocognitive and hormonal correlates of voluntary weight loss in humans," *Cell Metabolism* (2018). DOI: 10.1016/j.cmet.2018.09.024

## Provided by Cell Press

Citation: Weight loss success linked with active self-control regions of the brain (2018, October 18) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2018-10-weight-loss-success-linked-self-control.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.