

Yo-yoing weight, blood pressure, cholesterol and blood sugar readings may raise heart attack and stroke risk

October 1 2018

Fluctuations in weight, blood pressure, cholesterol and/or blood sugar levels in otherwise healthy people may be associated with a higher risk of heart attack, stroke and death from any cause compared to people with more stable readings, according to new research in the American Heart Association's journal *Circulation*.

This is the first study to suggest that high variability of these risk factors has a negative impact on relatively healthy people. The study is also the first to indicate that having multiple measures with high variability adds to the risk.

Compared to people who had stable measurements during an average 5.5 year follow-up period, those with the highest amount of variability (in the upper 25 percent) on all measurements were:

- 127 percent more likely to die;
- 43 percent more likely to have a [heart attack](#);
- 41 percent more likely to have a stroke.

Using data from the Korean National Health Insurance system, researchers examined data on 6,748,773 people who had no previous heart attacks and were free of diabetes, [high blood pressure](#), or [high cholesterol](#) at the beginning of the study. All participants had at least three health examinations between 2005 and 2012 (every two years is

recommended in the system). Records of the exams documented [body weight](#), fasting blood sugar, systolic (top number) blood pressure and total cholesterol.

Because high variability could result from either positive or negative changes, the researchers looked separately at the effect of variability in participants who were more than 5 percent improved or worsened on each measurement. In both the improved and the worsened groups, high variability was associated with a significantly higher risk of death.

"Healthcare providers should pay attention to the variability in measurements of a patient's [blood pressure](#), cholesterol and glucose levels as well as body weight. Trying to stabilize these measurements may be an important step in helping them improve their health," said Seung-Hwan Lee, M.D., Ph.D., senior author of the study and professor of endocrinology at the College of Medicine of the Catholic University of Korea in Seoul, South Korea.

The study was observational, which means that it cannot prove a cause-and-effect relationship between high variability and the risk of heart attacks, strokes or death from any cause. The study also did not delve into the reasons behind the fluctuations in the participants' risk factor measurements.

"It is not certain whether these results from Korea would apply to the United States. However, several previous studies on variability were performed in other populations, suggesting that it is likely to be a common phenomenon," Lee said.

Provided by American Heart Association

Citation: Yo-yoing weight, blood pressure, cholesterol and blood sugar readings may raise heart

attack and stroke risk (2018, October 1) retrieved 17 July 2024 from
<https://medicalxpress.com/news/2018-10-yo-yoing-weight-blood-pressure-cholesterol.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.