


Fewer than 5 in 100 Australian and NZ interventional cardiologists are women

November 30 2018, by Ivy Shih



Credit: AI-generated image ([disclaimer](#))

A new study has highlighted a critical gender gap in interventional cardiology, the medical field specialising in treating blocked arteries in heart disease.

There are only 19 female interventional cardiologists in all of Australia

and New Zealand – just 4.8 percent of that workforce, a new study by medical researchers, including a UNSW Ph.D. student, has shown.

In light of the findings, the researchers have called for change to increase much-needed diversity in the field – and to improve clinical care and outcomes for [women](#). With cardiovascular disease the main cause of death for both men and women, the study is relevant to a large number of Australians.

The findings, published in the *Journal of the American College of Cardiology*, are the result of research performed by Women in Interventional Cardiology of Australian and New Zealand (WiiCAN – pronounced 'we can').

On top of the severe gender disparity in interventional cardiology, the data also reveal that only 15 percent of all cardiologists across the two countries are women, that 73 percent of [public hospitals](#) have no female interventional cardiologists at all, and that three Australian states have none either.

One of the lead authors, interventional cardiologist Dr. Sonya Burgess, is working at Nepean Hospital in Sydney. She is also completing a doctorate at UNSW Sydney, studying patients with heart attacks and the importance of complete revascularisation (treatment) of the coronary arteries for these patients. She says there are many barriers for female trainees.

"I started my cardiology training in 2007 in New Zealand, the country that first gave women the vote, yet also a country that now has had more female prime ministers than female interventional cardiologists," she says.

"The reasons for this are complex, but can be overcome with positive

changes," Dr. Burgess says.

Their paper discusses factors contributing to this gap, including the importance of role models, the expectations of training and society, the impact of geography, occupational radiation, cardiology's workplace culture, unconscious gender bias, and strategies for promoting change.


The authors call for more evidence-based solutions to tackle the problem, including systematic approaches to actively address disparity and achieve more visibility, so trainees see female role models in leadership positions, in faculty at conferences and in training hospitals.

"We need leaders who believe the data is important and who value diversity, to allow us to change from a culture of de-facto exclusion to one of active inclusion," Dr. Burgess says.

One of the benefits of a more diverse workforce is better outcomes – for everyone.

Gender distribution in the medical workforce (2015-2017)

■ Female ■ Male



Source: Burgess, S. et al. J Am Coll Cardiol. 2018; 72(21):2663 - 7 • Get the data • Created with Datawrapper

Gender distribution in the medical workforce (2015-2017). Credit: Burgess, S. et al. J Am Coll Cardiol. 2018; 72(21):2663 - 7

Dr. Burgess says there is growing evidence showing that increased gender diversity in the profession can improve care and outcomes of female patients with heart disease.

"It is now well documented that women with [heart disease](#) receive less treatment, fewer appropriate medications, and experience higher mortality compared to men," she says.

"At the same time, several studies have reported better clinical outcomes for patients treated by female cardiologists and physicians, notably including improved survival rates for female [heart attack](#) patients when treated by a female doctor."

New social network part of strategy to address gender gap

To help address some of these issues and create a community of women, Dr. Burgess co-founded WiiCAN.

Dr. Burgess helped found the social network in 2017, when she met the other two co-founders, Dr. Sarah Zaman from Monash Medical Centre and Dr. Elizabeth Shaw from Hornsby Ku-ring-gai Hospital, at a conference that brought together Australian and New Zealand interventional cardiologists. Their goal is to give a voice to female cardiology trainees, patients and colleagues, so they would not feel isolated in a male-dominated workforce.

"When we met, it was incredibly special for us because it was essentially unheard of to have three female interventional cardiologists together in this country, we certainly had never experienced it," Dr. Burgess says.

"As a group of female interventional cardiologists, we were virtually

invisible. Visibility matters, especially at meetings like that, because they inspire us, and they are supposed to.

"We need to acknowledge that there are more than 19 women in Australia and New Zealand who have ever wanted to be interventional cardiologists, and there are certainly more than 19 women capable of it.

"We want to give all trainees a sense of belonging, and make sure they know that they could become successful leaders – in short, we need to actively facilitate change," Dr. Burgess says.

Women in Interventional Cardiology Practice in Australia

- 42 percent of doctors are female
- 36 percent of specialists are female
- 15 percent of cardiologists are female
- 4.8 percent of interventional cardiologists are female
- 73 percent of public hospitals have no female interventional cardiologists
- In Australia 3/8 states have no female interventional cardiologist

More information: Sonya Burgess et al. Women in Medicine, *Journal of the American College of Cardiology* (2018). [DOI: 10.1016/j.jacc.2018.08.2198](https://doi.org/10.1016/j.jacc.2018.08.2198)

Provided by University of New South Wales

Citation: Fewer than 5 in 100 Australian and NZ interventional cardiologists are women (2018, November 30) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-11-australian-nz-interventional-cardiologists-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.