


A new brain imaging study challenges the dominant theoretical model of autism spectrum disorders

November 15 2018



Credit: CEA

Autism spectrum disorders (ASDs) are neurodevelopmental disorders characterized by communication disorders, altered social interactions and sensory and behavioral abnormalities. Research in genetics and brain imaging suggests that abnormalities in the development of the brain concerning, in particular, the formation of neural networks and the functioning of synapses could be involved in the onset of ASD.

In recent years, neuroimaging studies have shown that people with ASDs have abnormalities in the functioning of certain brain areas known to be responsible for processing emotions, language and social skills. Research on the brain connectivity of people with ASD has shown in particular a

lack of "long-distance" connections and an increase in "short-distance" connectivity. These results formed the basis for the development of a theoretical model for understanding ASDs, according to which the lack of social attention and information processing observed (difficulty in understanding a situation as a whole, attention to certain details) is explained by a saturation of information processed by the brain linked to the increase in neural connectivity between adjacent brain areas.

However, Prof. Josselin Houenou, Professor of Psychiatry at UPEC, researcher at Inserm, practitioner at the Henri Mondor University Hospitals, AP-HP, and final author of the study published in *Brain*, explains: "The model is based on the study of heterogeneous pediatric populations, including autistic children of varying ages and symptomatology, and on non-specific neuroimaging methods which do not provide reliable measurement of short-distance connectivity."

In order to test the current model, the authors of the study used an innovation designed by Miguel Gevara, Jean-François Mangin and Cyril Poupon at NeuroSpin, namely an atlas specifically covering tractographic analysis of 63 "short-distance" connections from images obtained by diffusion MRI (dMRI). The dMRI demonstrates in vivo bundles of white matter in the brain by measuring the diffusion of water molecules, especially along axons. Tractography can then be used to reconstruct the paths of the nerve fiber bundles represented in the form of a tractogram.

The authors were thus able to study the links between "short-distance" connectivity and [social cognition](#) in a homogeneous adult population of persons with ASDs from the InFoR-Autism cohort* (27 persons with ASDs without intellectual disabilities and 31 control persons), which provides one of the richest databases per patient and control.

"The significance of the InFoR-Autism* cohort lies in the great richness

of the data collected for each subject included. We were thus able to link the neuroimaging results obtained with social cognition scores, measuring social ability, empathy, social motivation, etc.," recalls Dr. Marc-Antoine d'Albis, Henri Mondor Hospital, Inserm U955, first author of the study.

Discovery of a "short-distance" brain connectivity deficit associated with a lack of social interaction and empathy

These preliminary results are in stark contrast to the current theoretical model according to which the lack of social attention and information processing in people with ASDs is due to an increase in neural connectivity between adjacent brain areas. They now need to be confirmed by studies conducted in children with ASDs, as Prof. Houenou explains, "These results are preliminary but they suggest that these short-distance connectivity abnormalities may be involved in some of the social cognitive deficits present in autistic subjects. It is now necessary to conduct similar studies in children to confirm the results obtained in adults. Pediatric cohorts are necessary for studies in children of various ages—and therefore brain maturation—and this implies taking into account a much larger population of subjects.

If these initial conclusions are confirmed, it would make it possible to consider the development of new therapeutic approaches for deficits in social cognition. For example, transcranial magnetic stimulation could be explored because brain connectivity between adjacent areas is located on the surface of the [brain](#)."

More information: Marc-Antoine d'Albis et al. Local structural connectivity is associated with social cognition in autism spectrum disorder, *Brain* (2018). [DOI: 10.1093/brain/awy275](https://doi.org/10.1093/brain/awy275)

Provided by CEA

Citation: A new brain imaging study challenges the dominant theoretical model of autism spectrum disorders (2018, November 15) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-11-brain-imaging-dominant-theoretical-autism.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.