

Children's vaccinations and development checks prevent hospital admissions in childhood

November 5 2018

Children who receive nursery vaccinations and development checks are less likely to be admitted to hospital during childhood years.

This is the finding of the largest ever analysis examining how infant vaccinations and NHS [development](#) checks in primary care are linked to [children's](#) risk of emergency [hospital](#) admissions.

The study, by researchers from Imperial College London, tracked 319,730 children in England from birth for 13 years from 2000 to 2013.

The results, published in the journal *BMC Medicine*, suggest up to 13,500 children a year in England may be missing out on vaccines, and 83,000 may be missing development checks.

The researchers, from Imperial's School of Public Health, revealed infants who did not receive their vaccinations were twice as likely to be have an emergency hospital [admission](#) than children who received their vaccinations.

Young children normally have a one in four risk of being admitted to hospital in their first year of life—the research team revealed that for infants who do not receive their vaccinations, this risk increases to one in two.

The study, funded by the National Institute of Health Research, also found preschool children who did not complete their vaccinations were six times as likely to be admitted to hospital with a [vaccine](#)-preventable infectious illness such as measles.

Infants who missed out on [health](#) and development checks, in which general practitioners (GPs) or health visitors assess healthy mental and physical development—were over four and a half times as likely to end up in hospital.

The study also revealed that children whose vaccinations were delayed also had a higher risk of an emergency hospital admission in their childhood years.

Professor Sonia Saxena, lead author of the research from Imperial's School of Public Health, said: "Despite the wealth of evidence for vaccination in driving down serious illness and preventable deaths in children, the benefit of vaccines is still disputed in some sectors of society. This study shows getting your child vaccinated safeguards their future health against the risk of preventable hospital admission. And it prevents not just against infectious illness, but against conditions that can get worse when children are exposed to infection"

Professor Saxena, who established the Child Health Unit at Imperial, added: "This research also demonstrates how having a systematic preventive programme in the early years of life can protect the health of the population in later life, as well as reducing health costs from hospital admissions. It is a success marker for the NHS, and should encourage other nations to invest in universal health coverage. This would help achieve the UN Sustainable Development Goal 3—providing access to health care and vaccines for all by 2030."

Children in the UK receive vaccinations between the ages of 8 weeks

and 13 years old, protecting them against a range of illnesses including meningitis, measles and whooping cough.

Children are also invited for NHS health and development checks in their first few days of life, then at around two months old, one year old and two years old. These checks test for a number of things including any problems with growth, hearing and vision.

The research team, which followed children from a representative sample of 363 GP practices across England, found the most common reason for hospital admissions was an infectious illness, which was responsible for more than a third of admissions overall.

Children who had a chronic condition, such as asthma, diabetes or epilepsy, were at particularly high risk of an emergency hospital admission if they missed out on vaccinations or development checks. For instance, children with asthma had a fivefold risk if they missed checks and vaccinations; for children with epilepsy the risk was elevenfold.

The study revealed 98 per cent of all infants studied were fully vaccinated, and 87 per cent had received developmental checks. Further analysis revealed children who didn't receive vaccinations or developmental checks were more likely to live in deprived areas, have a teenage mother, or be first born.

Dr. Elizabeth Cecil, co-author from Imperial's Department of Primary Care and Public Health explained: "The fact that the majority of children received vaccinations and developmental checks is a fantastic achievement for our health system. However, the children who slip through the net are at significant risk of illness, and these risks accrue across childhood."

Co-author Dr. Dougal Hargreaves added: "Children's health in the UK

lags behind other Western European countries, and emergency hospital admissions for children have been steadily rising over recent years. Ensuring as many children as possible receive vaccinations and development checks may help combat this rise and improve the health of children and young people.'

More information: Elizabeth Cecil et al, Impact of preventive primary care on children's unplanned hospital admissions: a population-based birth cohort study of UK children 2000–2013, *BMC Medicine* (2018). DOI: [10.1186/s12916-018-1142-3](https://doi.org/10.1186/s12916-018-1142-3)

Provided by Imperial College London

Citation: Children's vaccinations and development checks prevent hospital admissions in childhood (2018, November 5) retrieved 3 May 2024 from <https://medicalxpress.com/news/2018-11-children-vaccinations-hospital-admissions-childhood.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--