

Chronic exposure to excess noise may increase risk for heart disease, stroke

November 5 2018



Credit: CC0 Public Domain

Exposure to environmental noise appears to increase the risk of heart attacks and strokes by fueling the activity of a brain region involved in stress response. This response in turn promotes blood vessel inflammation, according to preliminary research to be presented in Chicago at the American Heart Association's Scientific Sessions 2018.

The findings reveal that people with the highest levels of chronic [noise](#) exposure—such as highway and airport noise—had an increased risk of suffering [cardiovascular events](#) such as heart attacks and strokes, regardless of other risk factors known to increase [cardiovascular risk](#).

The results of the study offer much-needed insight into the biological mechanisms of the well-known, but poorly understood, interplay between cardiovascular disease and chronic noise exposure, researchers said.

"A growing body of research reveals an association between [ambient noise](#) and cardiovascular disease, but the physiological mechanisms behind it have remained unclear," said study author Azar Radfar, M.D., Ph.D., a research fellow at the Massachusetts General Hospital in Boston. "We believe our findings offer an important insight into the biology behind this phenomenon."

Researchers analyzed the association between noise exposure and major cardiovascular events, such as heart attacks and strokes, among 499 people (average age 56 years), who had simultaneous PET and CT scan imaging of their brains and blood vessels. Diagnostic validation was done in a subset of 281 subjects.

All participants were free of [cardiovascular illness](#) and cancer at baseline. Using those images, the scientists assessed the activity of the amygdala—an area of the brain involved in stress regulation and emotional responses, among other functions. To capture cardiovascular risk, the researchers examined the participants' medical records following the initial imaging studies. Of the 499 participants, 40 experienced a cardiovascular event (e.g., heart attack or stroke) in the five years following the initial testing.

To gauge noise exposure, the researchers used participants' home

addresses and derived noise level estimates from the Department of Transportation's Aviation and Highway Noise Map.

People with the highest levels of noise exposure had higher levels of amygdalar activity and more inflammation in their arteries. Notably, these people also had a greater than three-fold risk of suffering a [heart attack](#) or a stroke and other major cardiovascular events, compared with people who had lower levels of noise exposure. That risk remained elevated even after the researchers accounted for other cardiovascular and environmental risk factors, including air pollution, high cholesterol, smoking and diabetes.

Additional analysis revealed that high levels of amygdalar activity appears to unleash a pathway that fuels cardiac risk by driving blood vessel inflammation, a well-known risk factor for cardiovascular disease.

The researchers caution that more research is needed to determine whether reduction in noise exposure could meaningfully lower cardiovascular risk and reduce the number of cardiovascular events on a population-wide scale.

In the meantime, however, the new study findings should propel clinicians to consider chronic exposure to high levels of ambient noise as an independent risk factor for [cardiovascular disease](#).

"Patients and their physicians should consider chronic [noise exposure](#) when assessing cardiovascular risk and may wish to take steps to minimize or mitigate such chronic exposure," Radfar said.

Provided by American Heart Association

Citation: Chronic exposure to excess noise may increase risk for heart disease, stroke (2018,

November 5) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-11-chronic-exposure-excess-noise-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.