

## Drug overdose mortality rates highest in US

November 19 2018



(HealthDay)—In 2015, the United States had higher drug overdose mortality rates than those of many other Organisation for Economic Cooperation and Development member countries, according to a research



letter published online Nov. 13 in the Annals of Internal Medicine.

Yingxi Chen, M.D., Ph.D., from the National Cancer Institute in Rockville, Maryland, and colleagues compared trends and patterns in premature mortality rates due to <u>drug overdose</u> in 13 Organisation for Economic Co-operation and Development member countries.

The researchers found that the highest drug overdose mortality rates for both men and women were seen in the United States (35 and 20 deaths per 100,000 people, respectively), while Mexico had the lowest rates (1.0 and 0.2 deaths per 100,000 people, respectively). There was considerable variation in mortality trends by country. Estonia had the highest average annual percentage of change (AAPC) between 2001 and 2015 (AAPC, 6.9 percent in men and 7.9 percent in women), followed by the United States (AAPC, 4.3 and 5.3 percent, respectively); increases were also seen in Australia and the Netherlands. Despite high mortality rates, Norway had the largest decrease in drug overdose mortality (AAPC, -3.2 and -2.0 percent in men and women, respectively).

"Urgent public action is needed, as are improvements in maintaining the quality of systems and data," the authors write. "Detailed evaluations of the policies of countries with declining rates may help identify approaches that can be applied elsewhere to prevent further premature deaths."

**More information:** <u>Abstract/Full Text (subscription or payment may</u> <u>be required)</u>

Copyright © 2018 HealthDay. All rights reserved.

Citation: Drug overdose mortality rates highest in US (2018, November 19) retrieved 22 May



2024 from https://medicalxpress.com/news/2018-11-drug-overdose-mortality-highest.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.