

Drug resistant infections associated with higher in-hospital mortality rates in India

November 16 2018

In one of the largest studies to measure the burden of antibiotic resistance in a low- or middle-income country, researchers at the Center for Disease Dynamics, Economics & Policy report that in-hospital mortality is significantly higher among patients infected with multi-drug resistant (MDR) or extensively drug resistant (XDR) pathogens including *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae* and *Acinetobacter baumannii*.

Researchers analyzed antimicrobial susceptibility testing results and mortality outcomes for over 4,000 patients who visited one of ten tertiary or quaternary referral hospitals across India in 2015. Pathogens were classified as MDR or XDR based on drug susceptibility profiles proposed by the European and US Centers for Disease Control and Prevention. Mortality data was restricted to in-hospital deaths. Additional demographic and clinical data including age, sex, place of infection acquisition, and location in the hospital (i.e., intensive care unit [ICU] or non-intensive care unit) were also collected.

The overall mortality rate among all [study participants](#) was 13.1 percent, with mortality as high as 29.0 percent among patients infected with *A. baumannii*. Patients who died were more likely to have been older and admitted to the ICU at the time of testing. Researchers also found that among MDR infections, those caused by Gram-negative bacteria were associated with higher mortality rates compared to those caused by Gram-positive bacteria, with rates of 17.7 percent and 10.8 percent, respectively.

Study results indicate that patients who acquired MDR bacterial infections were 1.57 times more likely to die, compared to patients with similar susceptible infections, while patients who acquired XDR infections were 2.65 times more likely to die when accounting for age, sex, site of [infection](#), and the number of coinfections.

In both the ICU and non-ICU, odds of mortality were higher among patients with XDR infections; this association was driven by Gram-negative infections (e.g., XDR *K. pneumoniae*) highlighting the importance of rapidly identifying these infections among all [patients](#).

In India, MDR and XDR Gram-negative bacterial infections are frequent, and the availability of effective antibiotic therapies are declining. This study provides greater insight into the urgent need to increase surveillance, research, and antimicrobial stewardship efforts worldwide. The [researchers](#) further note that these findings on the mortality burden of antibiotic resistance can aid in the development of policy efforts to prioritize antibiotic resistance as a global public health threat and to inform future efforts to quantify and track the burden of resistance across low- and [middle-income countries](#).

More information: Sumanth Gandra et al, The mortality burden of multidrug-resistant pathogens in India: a retrospective observational study, *Clinical Infectious Diseases* (2018). [DOI: 10.1093/cid/ciy955](https://doi.org/10.1093/cid/ciy955)

Provided by Center for Disease Dynamics, Economics & Policy

Citation: Drug resistant infections associated with higher in-hospital mortality rates in India (2018, November 16) retrieved 13 March 2024 from <https://medicalxpress.com/news/2018-11-drug-resistant-infections-higher-in-hospital.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.