


Gut microbiota of mouse pups regulated by hydrogen peroxide produced through lactation

November 15 2018



Credit: CC0 Public Domain

A recent animal study published in *The FASEB Journal* explores the uncharted territory of how the gut microbiome develops in infancy. It was well known that breast-fed mouse pups have a simple bacterial

population dominated by the health-promoting *Lactobacillus* bacteria. But the molecular mechanism of how mouse mother's milk suppresses a more diverse gut microbiome has been unclear.

Previously, researchers examined the role of L-amino acid oxidase 1 (LAO1)—an enzyme that converts L-amino acids into keto acids, ammonia, and hydrogen peroxide (H₂O₂)—in mouse mothers' [milk](#). That study demonstrated that the H₂O₂ in milk prevented the mother's mammary gland from developing bacterial infectious disease, such as mastitis.

The same group of researchers conducted a more recent mouse study to consider whether the H₂O₂ generated by LAO1 might impact the development of infant gut microbiota via breast-feeding. Indeed, they found that LAO1 suppressed microbial diversity during the pups' gut microbiota development.

To conduct the study, researchers examined the amount of *Lactobacillus* in the microbiota of two groups of breast-fed infant mice: a control group and a group with a deleted (or "knockout") LAO1 gene. As expected, the mice in the control group displayed *Lactobacillus*-dominated microbiota. The microbiota of the LAO1-knockout mice, however, displayed lower levels of *Lactobacillus*. The study suggests for the first time that H₂O₂ acts as a gatekeeper for bacterial selection and the development of [gut microbiota](#) during lactation in mice.

"While there are many differences between human and [mouse](#) mothers' milk, both types of neonates have a low-diversity [microbiota](#), which might be important to reducing their risk of disease in adulthood," said Kentaro Nagaoka, Ph.D., an associate professor at the Laboratory of Veterinary Physiology within the Tokyo University of Agriculture and Technology's Department of Veterinary Medicine. "There is a need for

further research exploring how we can control breast milk compounds to ensure infants inherit the good microbiome."

"This was a clever and much-needed experiment, and the results have important human impacts," said Thoru Pederson, Ph.D., Editor-in-Chief of *The FASEB Journal*.

Provided by Federation of American Societies for Experimental Biology

Citation: Gut microbiota of mouse pups regulated by hydrogen peroxide produced through lactation (2018, November 15) retrieved 27 April 2024 from <https://medicalxpress.com/news/2018-11-gut-microbiota-mouse-pups-hydrogen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.