


Among heart attack survivors, drug reduces chances of second heart attack or stroke

November 7 2018



Credit: CC0 Public Domain

In a clinical trial involving 18,924 patients from 57 countries who had suffered a recent heart attack or threatened heart attack, researchers at the University of Colorado Anschutz Medical Campus and fellow scientists around the world have found that the cholesterol-lowering drug alirocumab reduced the chance of having additional heart problems or stroke.

The study was published today in the *New England Journal of Medicine*.

Alirocumab is in the class of drugs called PCSK9 antibodies.

"It works by increasing receptors on the liver that attract particles of LDL [cholesterol](#) from the blood and break them down. The result is that blood levels of LDL or 'bad' cholesterol decrease by approximately 50 percent, even when [patients](#) are already taking a statin," explained Gregory Schwartz MD, Ph.D., co-author of the study and professor of medicine at the University of Colorado School of Medicine.

The trial looked at patients who were at least 40-years-old, had been hospitalized with a heart attack or threatened heart attack (unstable angina), and had levels of LDL cholesterol of at least 70 mg per deciliter despite taking high doses of statins.

Half of the patients received alirocumab by self-injection under the skin every two weeks, and the other half received placebo injections. The patients were followed for an average of nearly three years. During that time, LDL cholesterol levels averaged 40 to 66 mg per deciliter in patients given alirocumab, compared with 93 to 103 mg per deciliter with placebo. Death from [coronary heart disease](#), another heart attack or episode of unstable angina, or a stroke occurred in 903 patients given alirocumab, compared with 1052 patients given the placebo, corresponding to a 15% reduction in risk.

"Statins have been the main cholesterol-lowering drugs for heart patients for more than 30 years, and they are very effective," Schwartz said.

"Now we know that we can improve the outcomes after a [heart attack](#) by adding alirocumab to statins in selected patients."

In the trial, alirocumab was safe and generally well-tolerated. The only common side effect with alirocumab was itching, redness, or swelling at

the injection site which was usually mild. It occurred in 3.8 percent of those given alirocumab, compared with 2.1 percent of patients who received the placebo.

Alirocumab was approved by the Food and Drug Administration in 2015 as a treatment for high cholesterol, but it has only now been shown to also reduce the risk of [heart](#) disease events and stroke.

Provided by CU Anschutz Medical Campus

Citation: Among heart attack survivors, drug reduces chances of second heart attack or stroke (2018, November 7) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-11-heart-survivors-drug-chances.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--