

Juice displaces milk and fruit in high school lunches

November 15 2018

High school students participating in school meal programs are less likely to select milk, whole fruit, and water when fruit juice is available, which on balance may decrease the nutritional quality of their lunches, according to a new study by the Rudd Center for Food Policy and Obesity at the University of Connecticut.

The National School Lunch program reaches over 30 million students, and while fruits, vegetables, and dairy served in the [school](#) lunch undoubtedly contribute to a healthy diet, the appropriate role of [juice](#) in children's diets has generated debate. Currently, the American Academy of Pediatrics recommends that children aged 7-18 consume no more than 8 ounces of juice daily, and juice is permitted to be served as part of the National School Lunch program only on certain days.

"Our findings suggest that on the days when juice is available as a choice in high school lunches, students do not select whole [fruit](#) and milk as often. This is a problem because compared to juice, milk and whole fruit are better sources of three nutrients of concern for adolescents—calcium, vitamin D, and fiber," says Marlene Schwartz, Director of the UConn Rudd Center and co-author of the study.

The study, published in the *Journal of Nutrition Education and Behavior*, utilized cafeteria register data from 3 low-income, Northeast high schools over one school year to determine whether [high school](#) students select different meal options on days when juice is available. The study also looked at the sales of a la carte beverages, which included both

water and 100% juice.

Key findings include:

- On juice days, 9.9% fewer milks and 7.4% fewer whole fruit servings were selected with lunches.
- 8.2% fewer bottles of water and 24.4% fewer bottles of 100% juice were sold a la carte when juice was offered.

"The potential nutritional impact of these substitutions is important to consider. For instance, an 8-ounce serving of apple juice has no vitamin D, 285 fewer grams of calcium, and 116 fewer grams of potassium compared to an 8-ounce serving of 1% milk," says Rebecca Boehm, lead author of the study and a University of Connecticut Postdoctoral Fellow with the Rudd Center for Food Policy and Obesity and the Zwick Center for Food and Resource Policy.

More information: Rebecca Boehm et al, Juice Displaces Milk and Fruit in High School Lunches, *Journal of Nutrition Education and Behavior* (2018). [DOI: 10.1016/j.jneb.2018.09.013](https://doi.org/10.1016/j.jneb.2018.09.013)

Provided by Elsevier

Citation: Juice displaces milk and fruit in high school lunches (2018, November 15) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-11-juice-displaces-fruit-high-school.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.