


Researchers identify a mechanism that fuels cancer cells' growth

November 14 2018, by Denise Heady



Credit: Min Yu (Eli and Edythe Broad Center for Regenerative Medicine and Stem Cell Research at USC), USC Norris Comprehensive Cancer Center

Scientists at the UCLA Jonsson Comprehensive Cancer Center have identified sodium glucose transporter 2, or SGLT2, as a mechanism that

lung cancer cells can utilize to obtain glucose, which is key to their survival and promotes tumor growth. The finding provides evidence that SGLT2 may be a novel biomarker that scientists can use to help diagnose precancerous lung lesions and early-stage lung cancers.

Cancer cells require large amounts of glucose to survive and grow. Past research has established that passive glucose transporters, or GLUTs, are the primary means the body uses to deliver glucose to tumors. However, more recent studies have found some cancers, such as prostate and pancreatic, also rely on SGLT2 to utilize glucose, which spurred the UCLA scientists to study the role of SGLT2 in relations to [lung cancer](#).

Finding ways to diagnose lung cancer earlier when it is easier to treat is imperative because it is currently the leading cause of cancer-related deaths in both men and women worldwide.

The researchers used [positron emission tomography](#), or PET scans, to capture SGLT activity in [lung cancer cells](#). They were able to use SGLT2 activity to monitor the effects of SGLT2 inhibitors to treat mice with genetic models of lung cancer and mice that had been implanted with human lung tumors.

The researchers found unusually high levels of SGLT2 in human specimens of lung cancer, showing that these tumors predominantly utilize SGLT2, but not GLUTs, to transport glucose in the early stages of [tumor growth](#) and even in pre-malignant lesions that precede the development of lung cancer.

The study suggests that lung cancer can be detected earlier, when it is much easier to treat. By using SGLT2 as a biomarker, lung cancer could not only be found earlier than it often is today, but it could potentially be discovered even before lesions become cancerous. The researchers also provided evidence that a common FDA-approved inhibitor drug (which

is currently used to treat diabetes) could help block SGLT2 activity in cancer cells. The inhibitor could be used to block [glucose uptake](#) and help reduce tumor growth.

The research is published in *Science Translational Medicine*.

More information: Claudio R. Scafoglio et al, Sodium-glucose transporter 2 is a diagnostic and therapeutic target for early-stage lung adenocarcinoma, *Science Translational Medicine* (2018). [DOI: 10.1126/scitranslmed.aat5933](#)

Provided by University of California, Los Angeles

Citation: Researchers identify a mechanism that fuels cancer cells' growth (2018, November 14) retrieved 19 April 2024 from <https://medicalxpress.com/news/2018-11-mechanism-fuels-cancer-cells-growth.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--